

Neuropsychology

Investigating the Interplay of Working Memory, Affective Symptoms, and Coping With Stress in Offspring of Parents With Huntington's Disease

Abigail E. Ciriegio, Anna C. Pfalzer, Lisa Hale, Katherine E. McDonell, Daniel O. Claassen, and Bruce E. Compas

Online First Publication, August 6, 2020. <http://dx.doi.org/10.1037/neu0000692>

CITATION

Ciriegio, A. E., Pfalzer, A. C., Hale, L., McDonell, K. E., Claassen, D. O., & Compas, B. E. (2020, August 6). Investigating the Interplay of Working Memory, Affective Symptoms, and Coping With Stress in Offspring of Parents With Huntington's Disease. *Neuropsychology*. Advance online publication. <http://dx.doi.org/10.1037/neu0000692>

Investigating the Interplay of Working Memory, Affective Symptoms, and Coping With Stress in Offspring of Parents With Huntington's Disease

Abigail E. Ciriegio
Vanderbilt University

Anna C. Pfalzer, Lisa Hale, Katherine E. McDonell,
and Daniel O. Claassen
Vanderbilt University Medical Center

Bruce E. Compas
Vanderbilt University






Objective: The offspring of parents with Huntington's disease (HD) are faced with substantial levels of chronic uncontrollable and unpredictable stress. These stressors may place them at heightened risk of psychological distress and negative effects on executive functioning. This study investigated working memory, secondary control coping strategies (e.g., cognitive reappraisal, acceptance, distraction), and symptoms of anxiety/depression in offspring at risk for HD. **Method:** Adolescent (ages 10–19) and young adult (ages 20–29) offspring ($n = 33$; mean [M] age = 19.12 years, standard deviation [SD] = 6.01 years; 61% female) of parents with HD were recruited in a Huntington Disease Society of America Level 1 Center of Excellence. Participants completed self-report measures of coping and neuropsychiatric symptoms (i.e., anxiety, depression) and a standardized working memory assessment. Pearson correlations and path analyses were used to test associations. **Results:** Participant scores on the working memory assessment were significantly lower compared with normative data, and scores on a mixed anxiety/depression scale revealed a significant elevation compared with normative data. Working memory, secondary control coping, and symptoms of anxiety/depression were significantly correlated. Analyses of the full model revealed that the total indirect effect of working memory on anxiety/depression through secondary control coping was significant ($\beta = -0.20$). **Conclusion:** Secondary control coping skills are an important factor in understanding the relationship between working memory and symptoms of anxiety/depression in the offspring of parents with HD. Future longitudinal research is needed to establish the direction of these associations. Working memory and coping skills represent potential targets for intervention to reduce the risk of anxiety/depression in this population.

Key Points

Question: How are working memory and coping with stress related to symptoms of anxiety and depression in adolescent and young adult offspring of parents with Huntington's disease (HD)?

Findings: Support was found for the indirect association of working memory with symptoms of anxiety and depression through secondary control coping (e.g., acceptance, cognitive reappraisal) in the offspring of parents with HD. **Importance:** Secondary control coping skills are an important factor in understanding the relationship between working memory and symptoms of anxiety and depression in the offspring of parents with HD. **Next Steps:** Future research should examine these associations using a longitudinal design and test working memory and skills for coping with stress as two potential targets for intervention that may reduce the risk of symptoms of anxiety and depression in the offspring of parents with HD.

Keywords: Huntington's disease, working memory, coping, anxiety, depression

 Abigail E. Ciriegio, Department of Psychology and Human Development, Vanderbilt University;  Anna C. Pfalzer, Lisa Hale,  Katherine E. McDonell, and  Daniel O. Claassen, Department of Neurology, Vanderbilt University Medical Center;  Bruce E. Compas, Department of Psychology and Human Development, Vanderbilt University.

Abigail E. Ciriegio served as lead for writing the original draft and writing review and editing and contributed equally to conceptualization and formal analysis. Anna C. Pfalzer served in a supporting role for formal analysis. Lisa Hale served in a supporting role for investigation. Katherine E. McDonell served in a supporting role for conceptualization and methodology. Daniel O. Claassen served as lead for conceptualization. Bruce E.

Compas served as lead for conceptualization and formal analysis. Abigail E. Ciriegio and Anna C. Pfalzer contributed to investigation equally. Abigail E. Ciriegio, Anna C. Pfalzer, Lisa Hale, and Daniel O. Claassen contributed to methodology equally. Daniel O. Claassen and Bruce E. Compas contributed to writing the original draft equally. Daniel O. Claassen and Bruce E. Compas contributed to writing review and editing equally.

Correspondence concerning this article should be addressed to Abigail E. Ciriegio or to Bruce E. Compas, Department of Psychology and Human Development, Vanderbilt University, Peabody 552, Nashville, TN 37203. E-mail: abigail.e.ciriegio@vanderbilt.edu or bruce.compas@vanderbilt.edu

Huntington's disease (HD) is an autosomal-dominant neurodegenerative disease caused by an expanded cytosine-adenine-guanine (CAG) repeat on the huntingtin (*HTT*) gene. Each offspring of a parent with HD has a 50% chance of carrying the genetic mutation, and all of those who inherit the genetic mutation will eventually develop the disease (Huntington's Disease Collaborative Research Group, 1993). Regardless of gene status, offspring are exposed to substantial levels of stress that heighten the risk for neuropsychiatric symptoms, especially anxiety and depression (Forrest Keenan, Miedzybrodzka, van Teijlingen, McKee, & Simpson, 2007; Vamos, Hambridge, Edwards, & Conaghan, 2007). Therefore, it is important to understand the skills that offspring need to cope with HD-related stress effectively and the neurocognitive resources that are needed to enact these coping skills. Previous research has identified coping skills that are important for offspring to cope with other types of parental disorders, including parental depression (e.g., Compas et al., 2010), and the importance of executive function skills that are associated with the ability to cope (Grech et al., 2016). However, these processes have not been assessed in the offspring of parents with HD, an especially vulnerable population. The current study focused on two factors uniquely important in understanding the impact of stress in the offspring of parents with HD: the ability to cope with stress and the proficiency of working memory, a type of executive function.

The offspring of parents with HD are faced with chronic, uncontrollable, and unpredictable stress, including observing the onset and progression of HD symptoms in a parent, caring for a parent, experiencing the ramifications of the neuropsychiatric symptoms of a parent, and realizing their own risk of developing HD (Forrest Keenan et al., 2007; Kavanaugh, 2014; Kjoelaas, Tillerås, & Feragen, 2020; Maxted, Simpson, & Weatherhead, 2014). Ultimately, much of this stress demands an ability to dynamically adapt to both static and fluctuating stressors. Coping strategies that focus on the ability to adapt to a problem, rather than trying to change a source of stress, are characterized as secondary control coping and include skills such as acceptance, cognitive reappraisal, positive thinking, and distraction (Compas et al., 2017; Connor-Smith, Compas, Wadsworth, Thomsen, & Saltzman, 2000). Secondary control coping skills are well suited to manage uncontrollable stress and may be particularly important for coping with HD-related stress (Downing, Williams, Leserman, & Paulsen, 2012). For example, the offspring of parents with HD may reappraise a parent's disease by focusing on the positive aspects of the time they have together or may identify ways that the disease has brought their family closer together. Prior research has shown that teaching secondary control coping skills as part of a cognitive-behavioral intervention can reduce the risk for anxiety and depression in adolescents of parents with major depressive disorder (e.g., Compas et al., 2010, 2015), suggesting that learning these skills may be a promising avenue for intervention with the offspring of parents with HD.

The use of complex cognitive coping strategies is predicated on intact and efficient executive function, including inhibitory control, cognitive flexibility, and working memory (Andreotti et al., 2013; Miyake & Friedman, 2012). For example, working memory skills require holding, updating, and manipulating information in a short time frame (e.g., recalling a sequence of numbers and reporting them in reverse order). Standardized tests of working memory measure the ability to manipulate information held in

short-term memory under "cold" conditions (i.e., nonemotional conditions). These skills can then be drawn upon to manipulate information (e.g., to think about a stressor in a different way through reappraisal) under conditions of stress and heightened emotional arousal (i.e., under "hot" conditions). For example, cognitive reappraisal, a type of secondary control coping, requires thinking about a stressful situation and simultaneously viewing it from an alternative perspective. Therefore, impairments in working memory may undermine the ability to utilize coping strategies that are most effective in the face of chronic, uncontrollable stress. Further, there is evidence that chronic stress directly impairs executive function skills and may indirectly impair the ability to use secondary control coping skills and other complex cognitive forms of coping (e.g., Liston, McEwen, & Casey, 2009).

Evidence for the association between working memory (and other aspects of executive function), secondary control coping skills, and emotional and behavioral problems has been established in other types of pediatric health populations (e.g., Campbell et al., 2009; Prussien et al., 2018; Robinson et al., 2015). For example, secondary control coping significantly mediated the relationship between a composite of standardized working memory assessment and total problem behaviors in a sample of adolescents with leukemia (Campbell et al., 2009). Similarly, a study using functional neuroimaging in survivors of childhood brain tumors found that increased brain activation in response to a working memory N-back task was associated with greater use of secondary control coping strategies and better psychosocial functioning (Robinson et al., 2015).

There is evidence of impairment in working memory skills in individuals with HD (Lahr et al., 2018), as well as in individuals with disorders of anxiety and depression (Moran, 2016; Snyder, 2013). Recent research has begun to examine brain development and cognitive function in individuals with juvenile HD and adolescent offspring at risk for HD (Lee et al., 2018; van der Plas et al., 2019). However, no studies, to our knowledge, have examined processes of coping and working memory in adolescent or young adult offspring of parents with HD. Understanding the possible impairments in working memory skills is especially important in offspring at risk for HD because problems with working memory may undermine the ability to efficiently cope with the stress presented by their parents' disease. Furthermore, an inability to manage levels of stress may ultimately place these offspring at greater risk for anxiety, depression, and other neuropsychiatric problems.

The present study investigated the associations among working memory, the use of secondary control coping strategies, and symptoms of anxiety and depression in a sample of adolescent and young adult offspring of parents with HD. The primary hypotheses were as follows: Offspring will have reduced working memory proficiency but higher levels of anxiety and depression compared with normative data (Hypothesis 1). Working memory, secondary control coping skills, and symptoms of anxiety and depression will be related at the bivariate level (Hypothesis 2). Specifically, working memory skills will be negatively associated with symptoms of anxiety and depression (Hypothesis 2a); working memory skills will be positively associated with secondary control coping skills (Hypothesis 2b); and secondary control coping skills will be negatively associated with symptoms of anxiety and depression (Hypothesis 2c). Finally, there will be an indirect association between

working memory and symptoms of anxiety and depression through secondary control coping (Hypothesis 3). Participants' age, levels of stress related to their parents' HD, and disease characteristics of their parents' HD (i.e., CAG repeat length) were controlled for in supplemental analyses. Given the high levels of stress in this population regardless of their genetic status, the ethical statutes preventing all offspring under the age of 18 from pursuing genetic testing (Huntington's Disease Society of America, 2009), and the fact that many young adult offspring choose not to be tested, it is important to study the relationship between working memory, coping, and symptoms of anxiety and depression in all offspring of parents with HD regardless of their gene status.

Method

Participants

Participants ($n = 33$) included 19 adolescent and 14 young adult offspring ages 10 to 29 years (mean [M] age = 19.12 years, standard deviation [SD] = 6.01 years; 61% female). The majority of the sample identified as White or Caucasian ($n = 32$), and only one participant identified as Black or African American. Parents with HD in the sample ranged in age from 32 to 58 years (M age = 45.58 years, $SD = 9.24$ years; 42% female) and had a mean CAG repeat length of 44.42 ($SD = 2.57$). All parents with HD in the sample identified as White or Caucasian. Although HD can affect individuals of all ethnic groups, it occurs much more frequently among those of European descent (Bates et al., 2015), which is representative of the current sample. Because of the ethical guidelines precluding individuals under age 18 years from undergoing genetic testing, the researchers were blind to the genetic status of all adolescents in the current sample. Of the individuals 18 years and older in the current sample who could pursue genetic testing, the researchers were only aware of the genetic status of a small subset ($n = 5$).

Procedure

Families were recruited through a Huntington Disease Society of America Level 1 Center of Excellence (COE) between October 2018 and December 2019. Eligibility requirements included (a) participants must use English as their primary language, (b) parents with HD must be part of the Huntington's Disease COE, (c) parents with HD may range in disease severity (i.e., presymptomatic, prodromal, manifest), (d) parents with HD must be over 30 years of age, and (e) parents with HD must have at least one offspring in the second 2 decades of life (10–19 [adolescent] or 20–29 [young adult]). Informed consent and assent were obtained from all parents and offspring prior to study enrollment and participation. This study was reviewed and approved by the Institutional Review Board. The medical director of the COE oversaw the recruitment of eligible families, and a member of the clinical team at the COE made the initial study introduction.

Measures

To assess demographics, coping strategies, and levels of neuropsychiatric symptoms in the sample, all participants were asked to

complete questionnaire data via REDCap, a secure online database (Harris et al., 2009).

Demographics. Demographic data for parents and offspring included information on age, sex, ethnic group, and education level. Disease characteristics (i.e., number of CAG repeats on the *HTT* gene) for parents with HD were extracted from medical records.

Stress and coping. The Responses to Stress Questionnaire–Huntington's Disease Version (RSQ-HD; Ciriegio et al., 2019; Connor-Smith et al., 2000) is a self-report measure that was used to identify sources of stress and coping strategies in response to stress from being in a family affected by HD. First, participants were provided with a list of 10 representative stressors related to HD (e.g., fear about my future, getting genetically tested, having to take on more caretaking responsibilities, feeling isolated or different from my peers) and asked to indicate if they had experienced any of these stressors in the previous 6 months. This portion of the RSQ-HD serves primarily as a cue to help respondents identify specific sources of stress that will be the focus of the subsequent coping items. A sum of the total number of stressor items endorsed was calculated for each participant. The second portion of the RSQ-HD includes 57 items reflecting coping and automatic responses to stressors, where each item is rated on a 4-point scale. Proportion scores are computed to identify five factors: primary control engagement coping, secondary control engagement coping, disengagement coping, involuntary engagement, and involuntary disengagement. Because of the uncontrollable nature of stress faced by the offspring of parents with HD, the current study focused on secondary control coping skills: *acceptance* (e.g., I realize I just have to live with things the way they are), *cognitive reappraisal* (e.g., I think about the things I'm learning from being from a family affected by HD, or that something good will come from it), *positive thinking* (e.g., I tell myself that I can get through this, or I will be okay), and *distraction* (e.g., I keep my mind off of the stressful parts of HD by doing something else). The RSQ-HD has demonstrated excellent internal consistency, test–retest reliability, and convergent and construct validity (Compas et al., 2017). Internal consistency for the secondary control coping scale in the current sample was $\alpha = .85$.

Anxiety and depression symptoms. Affective problems were assessed using the Youth Self Report (YSR; Achenbach & Rescorla, 2001) and the Adult Self Report (ASR; Achenbach & Rescorla, 2003) from the Achenbach System of Empirically Based Assessment (ASEBA). The ASEBA instruments are empirically driven, developmentally appropriate tools that assess psychopathology across the life span. These instruments have been developed and well validated in individuals ages 1½–90+ years, both in the United States and in international samples (Achenbach, Ivanova, & Rescorla, 2017). Offspring less than 18 years of age completed the YSR, and offspring 18 years of age and older completed the ASR. The YSR and ASR are self-report measures that yield both empirically based syndrome scales and scales oriented to the *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.; DSM-5; American Psychiatric Association, 2013). These measures have been shown to have excellent internal consistency, test–retest reliability, and construct validity. The normative samples for the YSR and ASR are representative of the U.S. population, providing adequate data on levels of emotional and behavioral problems in adolescents and young adults (Achenbach,

Dumenci, & Rescorla, 2002). The current study focused on the Mixed Anxious/Depressed Syndrome Scale as a broad indicator of affective problems, and this scale can be calculated in both the YSR and ASR. Ten items are included in both the child/adolescent and adult versions to reflect developmental continuities in some types of symptoms (e.g., sad, unhappy, depressed), but there are also various unique items in the child/adolescent and adult versions (three and eight items, respectively) to reflect developmental differences in other symptoms (e.g., afraid of going to school vs. poor relations with the opposite sex). The internal consistency for the Mixed Anxious/Depressed Syndrome Scale in the current sample was $\alpha = .84$ for the YSR and $\alpha = .90$ for the ASR.

Working memory skills. Working memory was assessed in the sample using the List Sorting Working Memory Task, a subtest of the National Institutes of Health (NIH) Toolbox Cognition Battery (Gershon et al., 2013; Tulsy et al., 2014). The List Sorting Working Memory Task requires the immediate recall and sequencing of various orally and visually presented stimuli. Pictures of different foods and animals are displayed with an accompanying audio recording (e.g., “elephant”), and the participant is asked to recall aloud, to the test administrator, the items in size order from smallest to biggest, first within a single dimension (either animals or food; 1 List) and then on two dimensions (foods and then animals; 2 List). A task score is calculated by summing the total number of items correctly recalled and sequenced on the 1 and 2 Lists. Raw scores can range from 0 to 26 and are converted to fully adjusted T scores ($M = 50$, $SD = 10$), which compares the individual’s score to the NIH Toolbox nationally representative normative sample while adjusting for age, sex, race, ethnicity, and educational attainment. This task has been shown to be developmentally appropriate and recommended for ages 7–85 years old (Tulsy et al., 2013, 2014). Administration of the NIH Toolbox Cognition Battery was completed electronically on an iPad and conducted by trained research assistants during in-person study visits at the COE.

Statistical Analyses

The mean T scores for working memory and symptoms of anxiety and depression in the current sample were compared to nationally representative normative T scores using independent-sample t tests. Bivariate associations between working memory, coping, and symptoms of anxiety and depression were identified using Pearson correlational analyses in Statistical Package for the

Social Sciences (SPSS; 26th ed.). Additionally, bivariate associations between working memory, coping, and symptoms of anxiety and depression with offspring age, sex, levels of stress (i.e., total sum of stressors on the RSQ-HD), and a disease characteristic of their parents’ HD (i.e., CAG repeats) were examined. Statistical power analyses indicated that with a sample of $n = 33$, there was adequate power ($\beta = .80$, $\alpha = .05$) to detect correlations of $r > .361$. To evaluate both the direct association between working memory and symptoms of anxiety and depression and the indirect association between working memory and symptoms of anxiety and depression accounted for by secondary control coping, Model 4 of the PROCESS macro (v3.4) for SPSS was used. The PROCESS macro is an ordinary least squares and logistic regression path-analysis and modeling tool that describes the total direct (path c , c') and indirect (path ab) effects through standardized regression coefficients and uses listwise deletion (Hayes, 2017). Current analyses were conducted with a 95% confidence interval (CI) for all effects. Offspring age, levels of stress, and CAG repeats of the offspring’s parent with HD were added as covariates in supplementary analyses of direct and indirect associations.

Results

The range, means, and standard deviations of measures are reported in Table 1. In support of the first hypothesis, adolescents and young adults obtained a mean T score of 46.36 ($SD = 9.71$) on the NIH Toolbox List Sorting Working Memory Task. This reflects a significant difference in working memory functioning compared with normative data, $t(32) = -2.2$, $p = .04$, and represents a small effect ($d = 0.36$). Also consistent with the first hypothesis, the sample’s mean T score on the Mixed Anxiety and Depression Scale on the YSR and ASR was 57.93 ($SD = 7.97$), revealing a large effect for elevated neuropsychiatric symptoms compared with normative data, $t(29) = 5.4$, $p < .001$, $d = 0.79$.

Support for the second hypothesis was found in bivariate correlation analyses (see Table 2). Working memory, as measured on the List Sorting Working Memory Task, was significantly correlated with secondary control coping on the RSQ-HD ($r = .40$, $p < .05$), and secondary control coping was significantly correlated with anxiety/depression on the YSR/ASR ($r = -.60$, $p < .01$). Further, there was an association between working memory and symptoms of anxiety/depression ($r = -.36$, $p = .051$) that was consistent with the hypothesis and approached significance. The level of stress, as measured by the total number of stressor items

Table 1

Descriptive Statistics for Measures of Offspring Working Memory, Coping, Symptoms of Anxiety and Depression, and Levels of Stress

Key constructs	Range	M	SD
NIH Toolbox List Sorting Working Memory Task	19–64	46.36	9.71
Secondary control coping	.12–.37	.26	.06
Mixed anxiety and depression	50–78	57.93	7.97
Levels of stress	0–9	2.84	2.16

Note. NIH = National Institutes of Health. Scores for the NIH Toolbox List Sorting Working Memory Task and the Mixed Anxiety and Depression Scale from the Youth Self Report (YSR)/Adult Self Report (ASR) are standardized T scores ($M = 50$, $SD = 10$). Secondary control coping scores, as measured on the Responses to Stress Questionnaire–Huntington’s Disease Version (RSQ-HD), are reported in ratio scores. Levels of stress is a sum of the total number of stressor items endorsed on the RSQ-HD (maximum score = 10).

Table 2
Correlations Between Offspring Age, Sex, Working Memory, Coping, Symptoms of Anxiety and Depression, Levels of Stress, and Parental CAG Repeat Length

Key constructs	1	2	3	4	5	6
1. Age	—					
2. Sex	.13	—				
3. NIH Toolbox List Sorting Working Memory Task	.02	-.27	—			
4. Secondary control coping	.31 [†]	.00	.40*	—		
5. Mixed anxiety and depression	-.15	.24	-.36 [†]	-.60**	—	
6. Levels of stress	-.02	.09	-.12	-.38*	.09	—
7. CAG repeat length in parents with HD	-.62**	-.20	-.06	-.20	-.18	.35 [†]

Note. CAG = cytosine-adenine-guanine; NIH = National Institutes of Health; HD = Huntington's disease.

[†] $p < .10$ (2 tailed). * $p < .05$ (2 tailed). ** $p < .01$ (2 tailed).

endorsed on the RSQ-HD, was significantly correlated with secondary control coping ($r = -.38, p = .03$). Parental CAG repeat length was significantly correlated with offspring age, $r = -.62, p < .001$. The sex of the offspring was not found to be significantly correlated with any measures.

Finally, the third hypothesis was confirmed (see Figure 1). Analyses of the full model using PROCESS revealed that the direct association between working memory and anxiety/depression (path c) was negative and approaching significance ($\beta = -.36, p = .051$). The direct association between working memory and secondary control coping (path a) was significant ($\beta = .38, p < .05$). In the full model, the effect of secondary control coping on anxiety/depression (path b) remained significant ($\beta = -.54, p < .01$), whereas the effect of working memory was no longer significant when coping was included in the model (path c'; $\beta = -.15$, not statistically significant [*ns*]). The total indirect effect of working memory on anxiety/depression through secondary control coping (path ab) was significant, $\beta = -.20$ (standard error [*SE*] = 0.09; 95% CI [-0.42, -0.06]).

Supplemental analyses were conducted in which offspring age, levels of HD-related stress in offspring, and CAG repeat length in parents with HD were included as covariates in the PROCESS analyses. The same pattern of significant paths was found when the age of the offspring was included in the model as a covariate, path a, $\beta = .34, p = .05$; path b, $\beta = -.58, p < .05$; path c, $\beta = -.35, p = .06$; path c', $\beta = -.15, ns$; path ab, $\beta = -.20$ (*SE* = 0.10; 95% CI [-0.44, -0.005]). Age was a significant

predictor of secondary control coping in the model ($\beta = .38, p = .03$) but was not associated with symptoms of anxiety and depression. In the model with levels of stress included as a covariate, the total indirect effect of working memory on symptoms of anxiety and depression through secondary control coping remained significant, path a, $\beta = .34, p = .06$; path b, $\beta = -.58, p < .05$; path c, $\beta = -.36, p = .06$; path c', $\beta = -.16, ns$; path ab, $\beta = -.19$ (*SE* = 0.10; 95% CI [-0.43, -0.02]). Level of stress was not a significant predictor of any measures in the full model. Furthermore, when parental CAG repeat length was included in the model as a covariate, the pattern of significant paths once again remained the same, path a, $\beta = .37, p = .05$; path b, $\beta = -.61, p < .05$; path c, $\beta = -.29, ns$; path c', $\beta = -.06, ns$; path ab, $\beta = -.23$ (*SE* = 0.12; 95% CI [-0.48, -0.02]).

Discussion

The offspring of parents with HD are faced with myriad chronic, uncontrollable, and unpredictable stressors related to the progression of the disease in a parent, as well as uncertainty and concerns about their own risk. Previous studies investigating other groups of adolescents and young adults faced with parental or familial illness have found that executive function skills and the skills used to cope with stress are important factors in understanding the association between stress and psychiatric symptoms (Prussien et al., 2018; Reising et al., 2018; Robinson et al., 2015). The current study extended this research to examine the potential role of working memory and secondary control coping skills as factors associated with symptoms of anxiety and depression in adolescents and young adults at risk for HD. Support was found for all three hypotheses. Most importantly, evidence was found for an indirect association of working memory with anxiety and depressive symptoms through secondary control coping.

Consistent with the first hypothesis, this sample of offspring of parents with HD scored significantly lower than the normative sample on a standardized measure of working memory and significantly higher than the normative sample on a standardized measure of symptoms of anxiety and depression. With regard to working memory, the observed level of problems was small in magnitude ($d = .36$). This finding is consistent with previous studies that have found variation in levels of cognitive function among gene-expanded and gene-nonexpanded offspring of parents with HD (Lee et al., 2018). It is noteworthy that the levels of problems in working memory in the current study were found in a




Figure 1. Direct and indirect associations of working memory and symptoms of anxiety and depression through secondary control coping without covariates. Standardized path coefficients, with standard errors in parentheses, are given. ⁺ $p < .10$. * $p < .05$.

sample of offspring of parents with HD that included both gene-positive and gene-negative individuals. The small effect may be due to the inclusion of both gene-expanded and gene-nonexpanded individuals in our sample. Furthermore, with regard to symptoms of anxiety and depression, the observed elevation in the current study was large in magnitude ($d = .79$). The findings from prior studies of affective symptoms in individuals at risk for and at various stages of HD have been mixed. For example, in a study of at-risk, prodromal, and symptomatic individuals with HD, Chisholm, Flavin, Paulsen, and Ready (2013) found that levels of depressive symptoms in individuals at risk for HD were comparable to those in a sample of healthy controls (Chisholm et al., 2013). In contrast, the current study used a measure of an empirically derived syndrome of mixed symptoms of anxiety and depression and compared the scores of the offspring of parents with HD with national normative data. The findings from the current study suggest that symptoms of anxiety and depression may be heightened in at-risk offspring of parents with HD as a consequence of the high levels of stress faced by these individuals.

Support was also found for the second hypothesis. Working memory skills were significantly positively associated with secondary control coping. This finding supports the hypothesis that working memory skills may be an important foundation for the ability to utilize complex coping strategies that are most effective in the face of chronic, uncontrollable stress. The current findings also supported the hypothesis that secondary control coping would be negatively associated with symptoms of anxiety/depression. Finally, the direct association between working memory and symptoms of anxiety and depression was negative, as hypothesized, and approached significance ($r = -.36, p = .051$). These findings indicate that adolescents and young adults who engaged more in the use of acceptance, cognitive reappraisal, and distraction when coping with HD-related stressors had fewer symptoms of anxiety and depression.

In support of the third and primary hypothesis, we found a significant indirect association between working memory and symptoms of anxiety and depression through secondary control coping. This suggests that secondary control coping is a unique and important factor in understanding the relationship between working memory and symptoms of anxiety and depression in the offspring of parents with HD. Furthermore, impairment in working memory may contribute to difficulties in using secondary control coping skills to deal with the stress of HD, which in turn is related to increased levels of anxiety and depression. These findings remained significant in supplemental analyses that included offspring age, levels of HD-related stress, and parental CAG repeat length as covariates. Taken together, these findings indicate two potential targets for cognitive and behavioral interventions that may be beneficial in aiding the psychological well-being of adolescent and young adult offspring of parents with HD: bolstering cognitive executive function skills and promoting the use of complex cognitive coping strategies, such as secondary control coping skills. These targets of intervention may be especially important because there are currently no available biologic interventions to slow the decline of functioning individuals at risk for HD.

The current study has several strengths. We employed a direct, performance-based measure of working memory from the NIH Toolbox Cognitive Battery, along with well-validated measures of secondary control coping skills (the RSQ-HD) and anxiety and

depression (the YSR/ASR), all with extensive normative data across age, sex, race, ethnicity, and education to allow for the interpretation of levels of these symptoms in adolescents and young adults. This sample included a wide age range of offspring of parents with HD spanning across adolescence and young adulthood.

We also note several limitations that can be addressed in future research studies. First, because of the small sample size, there was relatively low statistical power that could only detect medium to large effects, limiting the generalizability of the findings. Second, because it is cross-sectional, our study could not provide definitive conclusions about the directions of the associations among working memory, secondary control coping, and symptoms of anxiety and depression. A longitudinal design in which working memory is measured at an initial time point, followed by the measurement of coping at a second time point and symptoms of anxiety/depression at a third follow-up, could be used to establish temporal precedence and provide a more rigorous test of mediation. Third, the current findings are based on our full sample, which included both gene-positive and gene-negative offspring of parents with HD. However, none of the adolescents and only a small portion of the young adults had undergone genetic testing at the time of the study. Future research that includes genetic testing data on the offspring of parents with HD will be important to allow for comparison of those individuals with and without genetic mutations. Fourth, we examined an empirically derived syndrome of mixed anxiety/depression; future research could measure these symptoms separately to determine if there are effects that may be unique to anxiety as compared with depression. Fifth, we examined only one aspect of executive function, working memory, and future research should examine other domains of executive functioning, including inhibitory control and cognitive flexibility. Finally, randomized controlled trials to test interventions to enhance working memory skills (e.g., Jordan et al., 2020) or coping skills (e.g., Compas et al., 2010) or both (e.g., Bettis et al., 2017) would provide evidence for the role of these skills in reducing the risk for neuropsychiatric symptoms in the offspring of parents with HD.

In summary, the current study provides evidence for elevated symptoms of anxiety and depression in a sample of offspring of parents with HD. Levels of anxiety and depression were significantly associated with both difficulties in working memory and the use of complex cognitive coping skills. Further, we found evidence for an indirect path from working memory skills to the use of secondary control coping strategies, which in turn were related to lower levels of symptoms of anxiety and depression. These findings provide a basis for future longitudinal and intervention research in this population.

References

- Achenbach, T., Dumenci, L., & Rescorla, L. (2002). Ten-year comparisons of problems and competencies for national samples of youth: Self, parent, and teacher reports. *Journal of Emotional and Behavioral Disorders, 10*, 194–203. <http://dx.doi.org/10.1177/10634266020100040101>
- Achenbach, T., & Rescorla, L. (2001). *Manual for the ASEBA school-age forms and profiles*. Burlington, VT: University of Vermont, Vermont Center for Children, Youth & Families.

- Achenbach, T., & Rescorla, L. (2003). *Manual for the ASEBA adult forms and profiles*. Burlington, VT: University of Vermont, Vermont Center for Children, Youth & Families.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: Author.
- Achenbach, T. M., Ivanova, M. Y., & Rescorla, L. A. (2017). Empirically based assessment and taxonomy of psychopathology for ages 1½–90+ years: Developmental, multi-informant, and multicultural findings. *Comprehensive Psychiatry*, *79*, 4–18. <http://dx.doi.org/10.1016/j.comppsy.2017.03.006>
- Andreotti, C., Thigpen, J. E., Dunn, M. J., Watson, K., Potts, J., Reising, M. M., . . . Compas, B. E. (2013). Cognitive reappraisal and secondary control coping: Associations with working memory, positive and negative affect, and symptoms of anxiety/depression. *Anxiety, Stress & Coping: An International Journal*, *26*, 20–35. <http://dx.doi.org/10.1080/10615806.2011.631526>
- Bates, G. P., Dorsey, R., Gusella, J. F., Hayden, M. R., Kay, C., Leavitt, B. R., . . . Tabrizi, S. J. (2015). Huntington disease. *Nature Reviews Disease Primers*, *1*, 15005. <http://dx.doi.org/10.1038/nrdp.2015.5>
- Bettis, A. H., Coiro, M. J., England, J., Murphy, L. K., Zekowitz, R. L., DeJardins, L., . . . Compas, B. E. (2017). Comparison of two approaches to prevention of mental health problems in college students: Enhancing coping and executive function skills. *Journal of American College Health*, *65*, 313–322. <http://dx.doi.org/10.1080/07448481.2017.1312411>
- Campbell, L. K., Scaduto, M., Van Slyke, D., Niarhos, F., Whitlock, J. A., & Compas, B. E. (2009). Executive function, coping, and behavior in survivors of childhood acute lymphocytic leukemia. *Journal of Pediatric Psychology*, *34*, 317–327. <http://dx.doi.org/10.1093/jpepsy/jsp080>
- Chisholm, L. Z., Flavin, K. T., Paulsen, J. S., & Ready, R. (2013). Psychological well-being in persons affected by Huntington's disease: A comparison of at-risk, prodromal, and symptomatic groups. *Journal of Health Psychology*, *18*, 408–418. <http://dx.doi.org/10.1177/1359105312444646>
- Ciriegio, A., Hale, L., Pfalzer, A., McDonell, K., Riordan, H., Moroz, S., . . . Compas, B. (2019). *Understanding the effects of stress in families impacted by Huntington's disease: Associations of coping and psychiatric symptoms*. Presentation at the Annual Huntington Study Group Meeting, Sacramento, CA.
- Compas, B. E., Champion, J. E., Forehand, R., Cole, D. A., Reeslund, K. L., Fear, J., . . . Roberts, L. (2010). Coping and parenting: Mediators of 12-month outcomes of a family group cognitive-behavioral preventive intervention with families of depressed parents. *Journal of Consulting and Clinical Psychology*, *78*, 623–634. <http://dx.doi.org/10.1037/a0020459>
- Compas, B. E., Forehand, R., Thigpen, J., Hardcastle, E., Garai, E., McKee, L., . . . Sterba, S. (2015). Efficacy and moderators of a family group cognitive-behavioral preventive intervention for children of parents with depression. *Journal of Consulting and Clinical Psychology*, *83*, 541–553. <http://dx.doi.org/10.1037/a0039053>
- Compas, B. E., Jaser, S. S., Bettis, A. H., Watson, K. H., Gruhn, M. A., Dunbar, J. P., . . . Thigpen, J. C. (2017). Coping, emotion regulation, and psychopathology in childhood and adolescence: A meta-analysis and narrative review. *Psychological Bulletin*, *143*, 939–991. <http://dx.doi.org/10.1037/bul0000110>
- Connor-Smith, J. K., Compas, B. E., Wadsworth, M. E., Thomsen, A. H., & Saltzman, H. (2000). Responses to stress in adolescence: Measurement of coping and involuntary stress responses. *Journal of Consulting and Clinical Psychology*, *68*, 976–992. <http://dx.doi.org/10.1037/0022-006X.68.6.976>
- Downing, N. R., Williams, J. K., Leserman, A. L., & Paulsen, J. S. (2012). Couples' coping in prodromal Huntington disease: A mixed methods study. *Journal of Genetic Counseling*, *21*, 662–670. <http://dx.doi.org/10.1007/s10897-012-9480-3>
- Forrest Keenan, K., Miedzybrodzka, Z., van Teijlingen, E., McKee, L., & Simpson, S. A. (2007). Young people's experiences of growing up in a family affected by Huntington's disease. *Clinical Genetics*, *71*, 120–129. <http://dx.doi.org/10.1111/j.1399-0004.2006.00702.x>
- Gershon, R. C., Wagster, M. V., Hendrie, H. C., Fox, N. A., Cook, K. F., & Nowinski, C. J. (2013). NIH toolbox for assessment of neurological and behavioral function. *Neurology*, *80*, S2–S6. <http://dx.doi.org/10.1212/WNL.0b013e3182872e5f>
- Grech, L. B., Kiroopoulos, L. A., Kirby, K. M., Butler, E., Paine, M., & Hester, R. (2016). Coping mediates and moderates the relationship between executive functions and psychological adjustment in multiple sclerosis. *Neuropsychology*, *30*, 361–376. <http://dx.doi.org/10.1037/neu0000256>
- Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*, *42*, 377–381. <http://dx.doi.org/10.1016/j.jbi.2008.08.010>
- Hayes, A. F. (2017). *Introduction to mediation, moderation, and conditional process analysis* (2nd ed.). New York, NY: Guilford Press.
- Huntington's Disease Collaborative Research Group. (1993). A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell*, *72*, 971–983. [http://dx.doi.org/10.1016/0092-8674\(93\)90585-E](http://dx.doi.org/10.1016/0092-8674(93)90585-E)
- Huntington's Disease Society of America. (2009). *Genetic testing Huntington's disease: Family guide series*. New York, NY: Author.
- Jordan, L. C., Siciliano, R. E., Cole, D. A., Lee, C. A., Patel, N. J., Murphy, L. K., . . . Compas, B. E. (2020). Cognitive training in children with hypoplastic left heart syndrome: A pilot randomized trial. *Progress in Pediatric Cardiology*, *57*, 101185. <http://dx.doi.org/10.1016/j.ppedcard.2019.101185>
- Kavanaugh, M. (2014). Children and adolescents providing care to a parent with Huntington's disease: Disease symptoms, caregiving tasks and young carer well-being. *Child & Youth Care Forum*, *43*, 675–690. <http://dx.doi.org/10.1007/s10566-014-9258-x>
- Kjoelaas, S., Tillerås, K. H., & Feragen, K. B. (2020). The ripple effect: A qualitative overview of challenges when growing up in families affected by Huntington's disease. *Journal of Huntington's Disease*, *9*, 129–141. <http://dx.doi.org/10.3233/JHD-190377>
- Lahr, J., Minkova, L., Tabrizi, S. J., Stout, J. C., Klöppel, S., & Scheller, E., & the TrackOn-HD Investigators. (2018). Working memory-related effective connectivity in Huntington's disease patients. *Frontiers in Neurology*, *9*, 370. <http://dx.doi.org/10.3389/fneur.2018.00370>
- Lee, J. K., Conrad, A., Epping, E., Mathews, K., Magnotta, V., Dawson, J. D., & Nopoulos, P. (2018). Effect of trinucleotide repeats in the Huntington's gene on intelligence. *EBioMedicine*, *31*, 47–53. <http://dx.doi.org/10.1016/j.ebiom.2018.03.031>
- Liston, C., McEwen, B. S., & Casey, B. J. (2009). Psychosocial stress reversibly disrupts prefrontal processing and attentional control. *PNAS Proceedings of the National Academy of Sciences of the United States of America*, *106*, 912–917. <http://dx.doi.org/10.1073/pnas.0807041106>
- Maxted, C., Simpson, J., & Weatherhead, S. (2014). An exploration of the experience of Huntington's disease in family dyads: An interpretative phenomenological analysis. *Journal of Genetic Counseling*, *23*, 339–349. <http://dx.doi.org/10.1007/s10897-013-9666-3>
- Miyake, A., & Friedman, N. P. (2012). The nature and organization of individual differences in executive functions: Four general conclusions. *Current Directions in Psychological Science*, *21*, 8–14. <http://dx.doi.org/10.1177/0963721411429458>
- Moran, T. P. (2016). Anxiety and working memory capacity: A meta-analysis and narrative review. *Psychological Bulletin*, *142*, 831–864. <http://dx.doi.org/10.1037/bul0000051>
- Prussien, K. V., DeBaun, M. R., Yarboi, J., Bemis, H., McNally, C., Williams, E., & Compas, B. E. (2018). Cognitive function, coping, and

- depressive symptoms in children and adolescents with sickle cell disease. *Journal of Pediatric Psychology*, *43*, 543–551. <http://dx.doi.org/10.1093/jpepsy/jsx141>
- Reising, M. M., Bettis, A. H., Dunbar, J. P., Watson, K. H., Gruhn, M., Hoskinson, K. R., & Compas, B. E. (2018). Stress, coping, executive function, and brain activation in adolescent offspring of depressed and nondepressed mothers. *Child Neuropsychology*, *24*, 638–656. <http://dx.doi.org/10.1080/09297049.2017.1307950>
- Robinson, K. E., Pearson, M. M., Cannistraci, C. J., Anderson, A. W., Kuttesch, J. F., Jr., Wymer, K., . . . Compas, B. E. (2015). Functional neuroimaging of working memory in survivors of childhood brain tumors and healthy children: Associations with coping and psychosocial outcomes. *Child Neuropsychology*, *21*, 779–802. <http://dx.doi.org/10.1080/09297049.2014.924492>
- Snyder, H. R. (2013). Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: A meta-analysis and review. *Psychological Bulletin*, *139*, 81–132. <http://dx.doi.org/10.1037/a0028727>
- Tulsky, D. S., Carlozzi, N. E., Chevalier, N., Espy, K. A., Beaumont, J. L., & Mungas, D. (2013). V. NIH Toolbox Cognition Battery (CB): Measuring working memory. *Monographs of the Society for Research in Child Development*, *78*, 70–87. <http://dx.doi.org/10.1111/mono.12035>
- Tulsky, D. S., Carlozzi, N., Chiaravalloti, N. D., Beaumont, J. L., Kisala, P. A., Mungas, D., . . . Gershon, R. (2014). NIH Toolbox Cognition Battery (NIHTB-CB): List sorting test to measure working memory. *Journal of the International Neuropsychological Society*, *20*, 599–610. <http://dx.doi.org/10.1017/S135561771400040X>
- Vamos, M., Hambridge, J., Edwards, M., & Conaghan, J. (2007). The impact of Huntington's disease on family life. *Psychosomatics: Journal of Consultation and Liaison Psychiatry*, *48*, 400–404. <http://dx.doi.org/10.1176/appi.psy.48.5.400>
- van der Plas, E., Langbehn, D. R., Conrad, A. L., Kosciak, T. R., Tereshchenko, A., Epping, E. A., . . . Nopoulos, P. C. (2019). Abnormal brain development in child and adolescent carriers of mutant huntingtin. *Neurology*, *93*, e1021–e1030. <http://dx.doi.org/10.1212/WNL.00000000000008066>

Received March 27, 2020

Revision received June 9, 2020

Accepted June 10, 2020 ■