

# Stress and psychopathology in children and adolescents: is there evidence of specificity?

Susan D. McMahon,<sup>1</sup> Kathryn E. Grant,<sup>1</sup> Bruce E. Compas,<sup>2</sup> Audrey E. Thurm,<sup>1</sup> and Sydney Ey<sup>3</sup>

<sup>1</sup>DePaul University, USA; <sup>2</sup>University of Vermont, USA; <sup>3</sup>Pacific University, USA

Research on the relations between specific stressors and specific psychological outcomes among children and adolescents is reviewed. Specificity, the notion that particular risk factors are uniquely related to particular outcomes is discussed from a theoretical perspective, and models of specificity are described. Several domains of stressors are examined from a specificity framework (e.g., exposure to violence, abuse, and divorce/marital conflict) in relation to broad-band outcomes of internalizing and externalizing symptoms. Studies that tested for specificity conducted within the past 15 years are examined, and definitional problems are highlighted. Little evidence for specificity was found. Methodological problems in the literature and the lack of theory-driven specificity research are discussed, and directions for future research are identified. **Keywords:** Stress, psychopathology, specificity, child, adolescent.

Understanding the role of stress in the lives of children and adolescents is of both theoretical and practical significance. At the theoretical level, most prevailing models of developmental psychopathology recognize the potential importance of psychosocial stress in the etiology and maintenance of both internalizing and externalizing disorders in youth<sup>1</sup> (e.g., Cicchetti & Toth, 1991, 1997; Haggerty, Sherrod, Garmezy, & Rutter, 1994; Rutter, 1989). At the practical level, conditions and problems continue to worsen for children and adolescents. Levels of poverty, violence, and family adversity appear to be increasing (Children's Defense Fund, 1999), along with rates of emotional and behavioral problems in young people (Achenbach & Howell, 1993). Despite these trends, research on psychosocial stress and its relation to maladaptive outcomes in childhood and adolescence has lagged behind similar research with adults. Reviews concerning the impact of psychosocial stress on youth published in the past 15 years present a picture of a field early in its development (Cohen & Park, 1992; Compas, 1987; Johnson, 1986; Johnson & Bradlyn, 1988). In particular, research focused on testing theoretically-driven, developmentally-based models of *specific* relations between particular stressors and particular psychological outcomes has been lacking.

Specificity refers to the determination that a particular risk factor is uniquely related to a particular outcome (Garber & Hollon, 1991). In psychopathology research, a focus on specificity reflects an attempt to move beyond Selye's (1956) model in which 'general change' is associated with 'general distress'. Understanding specific pathways that link risk with maladjustment (e.g., death of a parent is linked to

depressive symptoms, but not disruptive behavior) is central to the goals of developmental psychopathology (Cicchetti & Cohen, 1995). Developmental psychopathology also considers equifinality, wherein many different avenues (of stressful experiences) can lead to the same outcome, and multifinality, in which similar conditions lead to multiple outcomes (Cicchetti & Toth, 1997). It is particularly important to examine specificity in children, as it is assumed that individual pathways evolve, and the causal processes contributing to these varied trajectories are multilevel. Developmental psychopathology acknowledges that many systems affect a child's behavior, including family variables, larger contextual variables, environmental stressors, and biological vulnerability. In this way, a model of developmental psychopathology does not presuppose evidence exists for specificity of environmental stressors and outcomes, although it does not disregard this possibility. It is only through theoretical guidance and empirical study that we may examine the strength of multifinality, equifinality and their equilibrium or middle ground, specificity.

In spite of the potential importance of this avenue of research, very few investigators have *intentionally* tested for specificity relations between particular stressors and particular psychological outcomes. Many studies have *explored* the relations between particular stressors and particular outcomes, yet to date there has been no attempt to integrate these findings. The purposes of the present review are to: 1) examine research on specificity relations between particular stressors and particular psychopathological outcomes, 2) summarize and integrate empirical findings, 3) define remaining concerns, and 4) highlight important directions for the next phase of research. The integration of these findings uses a lens of broad-band patterns of outcomes

<sup>1</sup> We use the term 'youth' throughout this review to refer to children and adolescents.

(i.e., internalizing versus externalizing behaviors). Although the field would benefit from moving toward a more narrow focus to capture the connections between specific stressors and specific outcomes, the variability in studies, particularly in terms of measurement and definition, and the lack of studies that focus on specificity, necessitate a more global review at this point in time.

### Methods of specificity research

The basic methodological requirement for specificity research is that the stressful experience is categorized by type of stressor, the psychological outcome is categorized by type of psychopathology, or both stressor and outcome are categorized. Although stressors can be categorized in numerous ways, categorization by type of event is the most appropriate. Categorizations based on frequency, duration, or severity of stress are less appropriate for specificity research, as specificity effects may be confounded with additive effects of stress (Rutter, 1989; Garmezy, 1987). For example, there is considerable evidence for the additive (and even multiplicative) effects of stress, such that exposure to more stressful life events or to more severe stress is predictive of more severe psychological symptoms (Rutter, 1989). However, much additional work needs to be done in conceptualizing and measuring stressors in a manner that captures the complexity of stressors in order to establish a meaningful examination of specificity in relation to psychological outcomes.

Categorization by type of stressful experience can be accomplished in several ways. Specific types of major life events may be compared. For example, parental divorce may be compared to death of a parent or child abuse. Major life events may also be subcategorized based on specific characteristics. For example, child abuse may be classified as physical abuse or sexual abuse (Manly, Cicchetti, & Barrett, 1994; Wolfe, Sas, & Wekerle, 1994). Exposure to violence may be classified as experienced versus witnessed violence, and chronic versus singular. One of the major challenges with this type of approach is that major events may subsume a large number of smaller events that have different characteristics. For example, divorce and exposure to violence may involve exposure to both conflict and loss. Another approach is to group individual stressors listed on cumulative measures of stressful life events based on theoretically derived categories. For example, events may be categorized as 'achievement events' versus 'affiliative events', or 'loss events' versus 'conflict events' (Sandler, Reynolds, Kliwer, & Ramirez, 1992). This type of categorization could be further refined to include various subcategories that describe stressors that are similar in nature in order to examine specificity between stressors and

outcomes. For example, in terms of loss events, death of a parent due to physical illness could be differentiated from death of a parent as a result of acts of violence. Unfortunately, there is no agreed upon taxonomy of event characteristics, and theoretical characteristics may be highly correlated with one another.

In order to understand the many complexities regarding understanding the influence of stressors on psychopathology, several steps need to be taken in order to develop a taxonomy of stressors for youth of various ages and backgrounds. Grant et al. (in press) suggest several steps to developing such a system. The first step would be to conduct structured interviews that assess events/circumstances as 'objectively threatening to the health or well-being of youth'. Then, lists of stressors can be generated and level of threat can be evaluated by external raters, taking into account the context of the stressor. Checklists can be developed for children of various ages and backgrounds, and norms can be established, using a method similar to Achenbach's method (1991; Achenbach and Dumenci, 2001) for examining psychopathology.

Psychological outcomes can be categorized by type using several different approaches. Single symptoms, clusters of symptoms (or syndromes), or disorders may be compared. For example, studies using the CBCL (Achenbach, 1991) may report results in terms of specific symptoms (e.g., endorsement of depressed mood vs. temper tantrums), narrow-band syndromes (e.g., anxious-depressed syndrome vs. aggressive syndrome), or broad-band syndromes (internalizing vs. externalizing syndromes). Categorization of psychological outcomes is complicated by the comorbidity among different outcomes. In addition, there are many methodological difficulties when stressors and psychopathology are examined together, including contamination of stress and outcomes, common method variance, and time precedence. Beyond categorization of stress and psychological outcome, specificity research does not require a specific methodology per se. However, design of a study directly affects the type and degree of specificity that may be gleaned from the results.

### Models of specificity

Three specificity designs have been utilized in the literature and are outlined in Figure 1: Stressor specific, Outcome specific, and Stressor-Outcome specific. The Stressor specific model includes several stressors and one outcome. This design allows for the identification of specificity of stressor in relation to a particular outcome, but does not allow for determination of specificity of outcome. In this model, the effect of each stressor on the outcome should be assessed independently of other stressors. The outcome could be identified as unique or common to various stressors. For example, Jarvelin, Moilanen,

*Stressor-Specific Model:*

Stressor #1  
 Stressor #2 → Specific psychopathology Outcome  
 Stressor #3

*Outcome-Specific Model:*

Psychopathology Outcome #1  
 Specific Stressor → Psychopathology Outcome #2  
 Psychopathology Outcome #3

*Stressor-Outcome Specific Model:*

Stressor #1            Psychopathology Outcome #1  
 Stressor #2    →    Psychopathology Outcome #2  
 Stressor #3            Psychopathology Outcome #3

**Figure 1** Models of specificity designs

Vikevainen-Tervonen, and Huttunen (1990) examined a number of stressful life events in relation to enuresis in children and found that divorce was the only stressful experience significantly related to enuresis. However, since only one psychological outcome was examined, it is unknown whether divorce would also predict other outcomes in this sample (e.g., depression, aggression).

The Outcome specific model includes several outcomes but only one stressor, allowing for the identification of specificity among outcomes but not among stressors. For example, Brody and Forehand (1990) examined rates of internalizing and externalizing symptoms among children exposed to marital conflict and found that internalizing symptoms were associated with parental conflict while externalizing symptoms were not. However, since only one stressful experience was examined, it is unknown whether other stressors would also predict high rates of internalizing symptoms relative to externalizing symptoms.


The Stressor-Outcome specific model includes a heterogeneous sample of stressors and a range of psychological outcomes, allowing for specificity of both stressor and outcome to be determined (Garber & Hollon, 1991). Each of the stressors can be examined in relation to each of the outcomes. Unlike Stressor specific and Outcome specific models, Stressor-Outcome specific models fully address whether a specific stressor is uniquely related to a specific outcome. Optimally, both the unique and common effects would be examined across different stressors. Sandler et al. (1992) utilized a Stressor-Outcome specific design to test the hypotheses that stressful events categorized as 'conflict events' would specifically predict externalizing symptoms and stressful events categorized as 'separation events' would specifically predict internalizing symptoms. Using multiple categories of stressful experiences

and multiple psychological outcomes, Sandler and colleagues (1992) found some evidence for their hypotheses (described further below).

Theoretically, models of specificity exist within the framework of broader conceptual models that include moderators and mediators of the relation between stress and psychological symptoms as well as reciprocal and dynamic relations among these variables (see Figure 2). Specific relations between particular stressors and particular outcomes may be mediated by psychological, biological, or social processes, or moderated by child characteristics or environmental context (Grant et al., in press). Thus, the specific form of symptoms may differ as a function of individual, family, or community factors, mediating processes, and the specific nature of the stressor. For example, death of a parent may be linked to depressive symptoms, and not disruptive behavior; but this relation may be particularly strong for older (adolescent) females, and may only exist when death of a parent is accompanied by additional household burdens. Very few studies testing full specificity designs (i.e., specific stressors linked to specific outcomes, via specific mediators, in the context of specific moderators) have been conducted. Although examination of specific mediators and moderators are beyond the scope of this study, it is important to acknowledge that they may influence our understanding of specific relations between stressors and outcomes. As a first step, however, it may be helpful to examine the results of available studies to take a closer look at the concept of specificity in relations between stressors and outcomes.

## Method of review

Literature was reviewed using both computer (PsychLit and PsychInfo) and manual methods (tracking citations). A computer-generated search was conducted using the following keywords: stress (or events or hassles) and psychopathology (or psychological symptoms or psychological disorder) and child (or adolescent), limited to empirical studies published in scientific journals in English. This search was repeated with specific additional stressors substituted for the stress term: abuse, divorce (or marital conflict), violence, poverty (or low income), illness (or death). Searches including these particular stressors were conducted because these stressors have not always been defined as stressors within specific studies but have received a great deal of research attention. Results of these additional searches were combined with the original search and duplicates removed, leaving approximately 1,500 original empirical articles on the relation between stress and psychological symptoms during childhood or adolescence published in scientific journals between 1987 and 2001. Research from this era was selected, because the last comprehensive reviews of



**Central Hypotheses of Conceptual Model:**

1. Stressors lead to psychopathology.
2. Moderators influence the relation between stressors and psychopathology.
3. Mediators explain the relation between stressors and psychopathology.
4. There is specificity in the relations among stressors, moderators, mediators, and psychopathology.
5. The relations among stressors, moderators, mediators, and psychopathology are reciprocal and dynamic.

**Figure 2** Conceptual model of the role of stress in the etiology of child/adolescent psychopathology (Grant et al., in press)

this literature appeared approximately 15 years ago (Compas, 1987; Johnson, 1986).

Approximately 15% of these studies (over 200 studies) tested for specific associations between particular stressors and particular outcomes (either cross-sectionally or longitudinally). Studies that defined stress as an environmental threat to the individual, or as a transactional relationship (*relationship* between person and environment *perceived* as taxing or exceeding resources and endangering well-being; Lazarus & Folkman, 1984), were included in this review. We excluded studies using a pure 'response-based' definition of stress (i.e., psychological or physiological distress in response to external events; Selye, 1974), as this definition is overly confounded with psychological symptoms.

The relative ease with which studies may examine specificity relations, without intentionally setting out to do so, has resulted in numerous studies with results regarding specificity effects. However, testing for specificity was not central to the research aims of most of these studies, and, in most cases, specific relations examined were not guided by developmental psychopathology theory. Rather, researchers included more than one stressor and more than one outcome in analyses focused on other research questions. Nonetheless, examination of results of these studies is an essential first step toward evaluating the validity of specificity theory.

The major focus of this review is on the more rigorous Stressor-Outcome specific studies (summarized in Table 1) and on the broad categorization of internalizing versus externalizing outcomes. In some cases, only internalizing or externalizing outcomes were examined, so these studies were not included in the review or the table. Given the preliminary nature of this review (it is the first to review evidence of

specificity effects), the dearth of theory-driven studies, and the variability in measurement of psychological outcomes in those studies that met inclusion criteria, examination of more specific psychological outcomes is not warranted (although many specific outcomes are reported in Table 1). Results of Outcome specific studies are also briefly reviewed (but not presented in the table), as determination of specificity of outcomes is most pertinent to the field of developmental psychopathology. Results of Stressor specific studies are not reviewed, as these studies do not allow us to contrast psychological outcomes based on the broad categorization system of internalizing versus externalizing outcomes.

Stressors that have been the focus of Stressor-Outcome specific studies include exposure to violence, abuse, and divorce/marital conflict.<sup>2</sup> In each of the areas, we looked for consistent patterns of specificity. That is, stressors within a particular category were related to internalizing but not externalizing outcomes, or externalizing but not internalizing outcomes. Mixed evidence for specificity was reported when there was not a consistent pattern of specificity (i.e., stressors were related to internalizing outcomes but other stressors within the same category were related to externalizing outcomes). No evidence of specificity was reported when particular stressors were not related to either externalizing or internalizing outcomes or were related to both internalizing and externalizing outcomes. Results of Stressor-Outcome and Outcome Specific studies are reviewed below, and findings of specificity for Stressor-Outcome studies (specificity, mixed specificity, no specificity) are presented in Table 1.

<sup>2</sup> Some of the studies fit into more than one category, so they are cross-referenced in Table 1.

**Table 1** Studies using 'Stressor-Outcome' designs to test for specificity between stress and symptoms (internalizing and externalizing) among children and adolescents

**Exposure to Violence**

Author	Sample	Design	Stress measure	Outcome measure	Findings
Attar, Guerra, & Tolan (1994)	384 children (1st, 2nd & 4th grade; 50% F; 57% AA, 43% L)	C/S & L (2 ts, 1 year apart)	self-report cumulative stress scale (SI)	teacher-report aggressive & anxious-depressed symptoms (TRF)(int & ext)	<b>Specificity for ext (not int):</b> transitions & exposure to violence (but not circumscribed events) predicted aggressive symptoms cross-sectionally (not anxious-depressed sx) & circumscribed events & exposure to violence (but not transitions) predicted aggressive symptoms over time
Durant, Pendergrast, & Cadenhead (1994)	224 adolescents (11-19 yrs; 56% F; 100% AA)	C/S	several types of exposure to violence (modified SECV), exposure to domestic violence (CTS)	self-report depressive sx (CDI), conduct symptoms (gang fighting, physical fights, domestic violence) (DYSSRDQ) (int & ext)	<b>Specificity for ext (not int):</b> exposure to violence & victimization by violence was correlated with all types of conduct symptoms; exposure to domestic violence was specifically related to domestic fighting
Gorman-Smith & Tolan (1998)	245 youth (11-15 yrs at Time 1; 100%M; 100% AA & L) & caregivers	L (2 ts, 1 year apart)	exposure to violence; stressful life events, family relations (dev by au), parenting (modified PYS)	parent, youth, & teacher report of anxiety/depression & aggression (int & ext)	<b>Mixed Specificity:</b> Exposure to violence was related to aggression & anxiety/depression, but other stressors were related to anxiety/depression, not aggression at Time 2
Levendosky & Graham-Bermann (1998)	121 children (7-12 yrs; 51% F); 60 of battered women & 61 controls (53% W, 47% AA)	C/S	witnessed battering of mother (CTS), witnessed psychological abuse of mother (VAWS); mother-report income	mother-report (CBCL) (int & ext)	<b>Specificity for int (not ext):</b> exposure to battering of mother was associated with internalizing (but not externalizing) symptoms; Exposure to psychological abuse of mother was associated with internalizing & externalizing symptoms
Lynch & Cicchetti (1998)	322 children (7-12 yrs): 188 maltreated, 134 nonmaltreated (37% F; 62% AA, 12% L, 2% O)	L (2 ts, 1 year apart)	exposure to violence (CVS), maltreatment (DSS records)	teacher report int & ext; child report PTSD sx (Levonn Measure), depression & anxiety (Checklist of Child Distress Symptoms) (int & ext)	<b>Mixed Specificity for exposure to violence:</b> exposure to violence was related to ext & depressive symptoms; Exposure to violence was specifically related to depressive symptoms (but not general ext or int)
McCloskey, Figueredo, & Koss (1995)	365 children (6-12 yrs; 50% F; 53% W, 35% L, 6% AA, 2% O) in shelters or treatment	C/S	exposure to domestic violence (e.g., verbal, physical, sexual, partner to mother, partner to child, mother to child)	parent & self-report psychological disorders (CBCL, CAS) (int & ext)	<b>Specificity for ext for sexual abuse:</b> Sexual abuse was specifically related to externalizing behavior (relationship); <b>No Specificity for maltreatment:</b> related to both int & ext behavior
					<b>Mixed Specificity:</b> exposure to domestic violence was associated with aggression, hyperactivity, delinquency, depressive symptoms, oppositional behavior, anxiety, separation anxiety, somatic & withdrawal symptoms (but not enuresis & encopresis)

Table 1 Continued

Exposure to Violence

Author	Sample	Design	Stress measure	Outcome measure	Findings
O'Keefe (1994a)	185 children (7–13 yrs; 49% F; 37% L, 21% AA, 42% W) in shelters	C/S	marital violence, mother-child aggression, father-child aggression (modified CTS)	parent report (CBCL) (int & ext)	<b>Mixed Specificity:</b> marital violence & mother-child aggression were both related to both internalizing & externalizing problems, but father-child aggression was not related to either internalizing or externalizing
O'Keefe (1997)	935 adolescents (14–20 yrs; 59% F; 53% L, 13% AA, 20% W, 7% AS, 7% O)	C/S	self-report inter-parental aggression, child physical abuse (modified CTS)	self-report (YSR) (int & ext)	<b>No Specificity:</b> both interparental aggression & physical abuse were related to both internalizing & externalizing behavior
Osofsky, Wewers, Hann, & Fick (1993)	53 children (9–12 yrs; 100% AA; NFDLP)	C/S	parent-report exposure to violence (SECV); family conflict (CTS)	parent report (SCSS, CBCL) (int & ext)	<b>Specificity for int (not ext):</b> witnessing violence & victimization were specifically related to internalizing syndrome, but not externalizing; no specificity found for family conflict (related to both int and ext)
Rossmann, Bingham, & Emde (1997)	86 children: 30 parent violence (4–9 yrs; 47% F; 44% W, 32% AA, 24% O), 14 dog attack (29% F; 47% W, 13% AA, 13% L, 27% O) & 42 mild stress controls (60% F; 90% W, 2% L, 8% O)	C/S	maternal report parental violence (CTS); dog attack; parent & child report mild stressors (modified LEQ)	parent-report PTSD (CBCL-PTSD, CDC, modified PTSD-RI) & int & ext sx (CBCL); self-report PTSD sx (PTSD-RI & CDC) (int & ext)	<b>Specificity for int (not ext):</b> parental violence group exhibited heightened rates of parent-report internalizing symptoms (CBCL-PTSD) relative to other 2 groups & higher rates on of PTSD sx (PTSD-RI) (& dog attack group higher) than mild stress group; dog attack group exhibited higher rates of child-report dissociative sx than children in the other 2 groups;
Sternberg, Lamb, Greenbaum et al. (1993)	110 children (8–12 yrs): 16 witnessed SV, 33 physically abused, 30 witnessed & abused, 31 controls (44%F; W)	C/S	exposure to interparental violence &/or physical abuse (DFS reports)	self-report depressive sx (CDI); int & ext syndromes (YSR), parent-report (CBCL) (int & ext)	<b>Specificity for ext (not int):</b> witnessing marital violence was associated with externalizing behaviors (no specificity for physical abuse)(parent report) (no specificity by child report)
Tang (1997)	59 children (6–13 yrs): 21 in shelters, 20 in tx, 18 intact families (56% F; 100% Chinese)	C/S	inter-parental verbal & physical abuse (mother report) or child abuse (verbal or physical) (modified CTS)	self-report of anxiety (STAI-C), depression (DSRS) & parent-report of aggression (CBCL) (int & ext)	<b>Mixed Specificity:</b> husband to wife physical abuse was associated with depression, anxiety & aggressive symptoms; interparental verbal & physical abuse was associated with both depression & anxiety symptoms; husband to wife verbal abuse & husband to child physical abuse were associated with depression & aggression symptoms; husband to child verbal abuse was associated with depression symptoms; mother to child physical abuse was associated with aggression symptoms

See also Jouriles, Barling, & O'Leary (1987) in Abuse. See also Jouriles et al., (1996a, b); Marttunen et al., (1994) in Divorce/Marital Conflict.

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
<b>Abuse</b> Ackerman, Newston, McPherson et al. (1998)	170 children (7–13 yrs): 127 sexually abused, 43 physically abused (49% F; 72% W, 28% AA)	C/S	physical abuse & sexual abuse; referred by hospitals & agencies; DHS reports	self & parent report 13 diagnoses (DICA); parent & teacher report (CBCL, TRF) (int & ext)	<b>Specificity for physical abuse ext</b> ( <b>not int</b> ): <i>physical abuse</i> was specifically related to <i>ODD &amp; Conduct Disorder</i> (DICA); physical abuse was associated with more behavior problems than sexual abuse (CBCL) <b>No Specificity for sexual abuse</b>
Bayatpour, Wells et al., (1992)	352 pregnant teenagers (12–19 yrs; 63% L, 19% W, 14% AA, 5% AS)	C/S	self-report physical abuse or sexual abuse (clinical interview)	self-report substance use, suicidal ideation & attempts (clinical interview) (int & ext)	<b>No Specificity for physical or sexual abuse</b> : Both physical abuse & sexual abuse were significantly associated with suicidality & substance use in pregnant teenagers
Boney-McCoy & Finkelhor (1995)	2000 children & adoles- cents (10–16 yrs; 48% F; 80%W, 10% AA, 7% H, 3% O)	C/S	self-report of sexual assault, violence by parents, non-parental family violence, simple assault, aggravated assault attempted kidnapping (dev by au)	self-report symptoms of PTSD (SC-90-R, modified by Sanders, Arata & Kilpatrick), sadness, & acting out with teacher (dev by au)(int & ext)	<b>Mixed Specificity for physical abuse</b> : <i>aggravated assault</i> specificity relating to <i>PTSD &amp; acting out at school</i> (but not sadness); <i>attempted kidnapping</i> was specifically related to <i>PTSD</i> ; <i>genitalia</i> <i>violence</i> was specifically related to <i>PTSD</i> & <i>acting out at school</i> (but not sadness)
Caviola & Schiff (1988)	270 adolescents (13–18 yrs): 150 abused (51%F); 60 chemically dependent (39%F); 60 controls (53%F); NFDP	C/S	sexual abuse, physical abuse, incest, & physical abuse & incest (treatment program & child abuse reporting records)	sx of suicide, delinquency, sexual acting out (method not described)(int & ext)	<b>No Specificity for sexual assault</b> <i>Specificity findings by gender</i> : simple assault relating specifically to sadness & acting out (but not PTSD)(boys); parental violence was associated with PTSD & acting out (but not sadness) (girls); non- parental family violence was specifically related to acting out at school (girls); attempted kidnapping related to sadness in girls & acting out in boys
Cohen, Spirito, Sterling et al., (1996)	105 abused children (12–18 yrs; 69% F; NFDP); inpatient psychiatric unit	C/S	physical abuse (by adult family member), sexual abuse (sexually coerced contact by adult) & boy sexual & physical abuse (based on admission)	self-report depressive sx (CDI), behavior problems (YSR), personality disorder, depression, conduct dis- order, substance abuse & psychosis (DICA-R & clinical eval)(int & ext)	<b>No Specificity for physical or sexual abuse</b> : for diagnoses based on type of abuse

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Crittenden, Claussen, & Sugarman (1994)	100 maltreated children (6–17 yrs; 47% F; 18% W, 37% AA, 33% L, 10% Haitian, 2% O)	C/S	physical abuse, physical neglect, both, risk for abuse & psychological maltreatment (standardized assessments, rating scales & interviews)	parent & case coordinator report of sx of conduct disorder, aggression, attention, anxiety-withdrawal, psychosis, motor excess (RBPC); self-report depressive sx (CDI)(int & ext)	<b>Specificity for int (not ext) for physical abuse:</b> <i>physical abuse &amp; depressive symptoms specifically related; physical neglect specifically related to inattention, anxiety-withdrawal, motor excess; emotional abuse specifically related to socialized aggression, inattention, anxiety withdrawal, psychotic symptoms; emotional neglect specifically related to inattention symptoms</i> <b>No Specificity for neglect</b>
DePaul & Arruabarrena (1995)	66 children (5–11 yrs): 17 physically abused, 24 physically neglected, 25 non-abused (100% Basque; NFDPP)	C/S	parent report physical abuse and physical neglect (Child Abuse Potential Inventory (CAP))	teacher report of sx (TRF) (int & ext)	<b>Specificity for int (not ext) for physical abuse:</b> <i>physical abuse was specifically related to internalizing symptoms</i> <b>Specificity for ext (not int) for neglect:</b> <i>physical neglect was specifically related to externalizing symptoms, compared to control group</i>
Deblinger, McLeer, Atkins, Ralph, & Foa (1989)	87 children (3–13 yrs): 29 sexually abused, 29 physically abused, 29 non-abused inpatient (47% F; NFDPP)	C/S	sexual abuse, physical abuse; hospital medical records	clinician reported sx of PTSD (re-experiencing phenomena, sexual acting out, avoidance behavior, & hyperarousal), suicidal ideation, somatic complaints (dev by au) (int only)	<b>Specificity for sexual abuse and sexual acting out:</b> <i>sexual abuse (not physical abuse) found specifically related to re-experiencing symptoms of PTSD (e.g. sexual acting out, sexually inappropriate behavior, avoidance)</i>
Dykman et al., (1997)	125 children (8–12 yrs): 109 abused in treatment & 16 controls (57% F; 67% W, 33% AA)	C/S	sexual abuse & physical abuse (psychiatric facilities & government agencies for abuse)	parent report PTSD (DICA), int & ext syndromes, sexually aggressive behavior (CBCL, DICA), teacher report (TRF), child report PTSD (DICA) (int & ext)	<b>Specificity for sexual abuse int (not ext, or sexual acting out):</b> <i>sexual abuse (not physical abuse) was specifically associated with PTSD and not other outcomes</i> <b>No Specificity for physical abuse</b>
Feldman et al., (1995)	193 children (8–12 yrs): 106 physically abused, 87 controls (29% F; 56% AA, 38% H, 5% W, 1% AS)	C/S	physical abuse (agency report), witnessing spouse/partner abuse (interview dev by au)	teacher report & parent report int & ext syndromes (CBCL, TRF), peer report of acceptability (dev by au) (int & ext)	<b>Specificity for ext (not int) for physical abuse:</b> <i>physical abuse (&amp; not witnessing spousal violence) was specifically associated with externalizing</i>



Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Friedrich, Jaworski, Huxsahl, & Bengtson (1997)	350 children (7–18 yrs): 72 inpatient sexually abused (80%F), 38 inpatient, likely abused (68% F), 165 inpatient control (42% F), 75 control (56% F; 'most W')	C/S	history of sexual abuse, physical abuse	self-report sx of dissociation (TSC-C), sexual concerns (R, TSC-C, CBCL); parent report MMPI, CDC, C-SBI) (int & ext)	<b>Specificity for sexual abuse int (not ext) &amp; sexual acting out:</b> <i>sexual abuse</i> was specifically related to <i>sex items</i> on the CBCL (parent report), but not by child report; no specificity found for sexual concerns (no other variables predicted it), some specificity was found for dissociation symptoms; duration & nature of sexual abuse (but not physical abuse) predicted dissociation
Gale, Thompson, Moran, & Sack (1988)	202 children (0–7 yrs; 90% W, 10% O): 37 sexually abused (76% F), 35 physically abused (51% F), 130 nonabused (42% F)	C/S	sexual abuse, physical abuse (referral from DSS & clinical evaluation)	therapist report sx of depression, anxiety, withdrawal, noncompliance, aggression, inappropriate sexual behavior, affective disorder, somatic complaints, suicidal ideation (int & ext)	<b>No Specificity for physical abuse</b> <b>Specificity for sexual abuse int (not ext) &amp; sexual acting out:</b> <i>sexual abuse</i> was specifically related to <i>sexual behavior problems</i> (compared to physical abuse & no abuse), & <i>withdrawal symptoms</i> (compared to no abuse); sexual abuse was also found to be specifically related to <i>lowered noncompliance</i>
Hernandez (1995)	6224 children (6th, 9th & 12th graders; 48% F; 85% W, 8% AI, 7% AA)	C/S	self-report physical abuse, incest, or extrafamilial sexual abuse (interview dev by au)	self-report eating disorder (2 of: binge eating, using laxatives or vomiting), anxiety sx, suicidal thoughts, substance use (dev by author)(int & ext)	<b>No Specificity for physical abuse</b> <b>No Specificity for physical or sexual abuse:</b> all three types of abuse were associated with eating disorders, anxiety, suicidal thoughts, substance use
Hernandez, Lodico, & DiClemente (1993)	2983 adolescents (9th & 12th grade; 100%M; 91% W, 9% AA) (412 abused)	C/S	self-reported incest, non-incest sexual abuse, physical (dev by au)	self-reported substance abuse, suicidal & delinquency sx (sexual aggressiveness, violent acts) (dev by au) (int & ext)	<b>Specificity for sexual abuse ext (not int or sexual acting out):</b> <i>sexual abuse</i> (both incest & extrafamilial) was specifically related to <i>drinking problems</i> (compared to physical abuse); no specificity for sexual aggression (high in both abuse groups) or for other delinquency behaviors (high in both groups except for a specific relation between <i>sexual abuse &amp; truancy</i> )
Jouriles, Barling, & O'Leary (1987)	45 children (5–13 yrs; 52% F) from maritally violent homes	C/S	parent report, witnessing inter-spousal aggression, parent-child aggression (CTS)	symptoms of conduct, depression, inattention, anxiety-withdrawal, psychosis, motor excess & socialized aggression (parent report, R-BPC) (int & ext)	<b>No Specificity for physical abuse</b> <b>Mixed Specificity for physical abuse:</b> <i>parent-child aggression</i> was related to <i>conduct problems, attention problems, anxiety-withdrawal, &amp; motor excess in boys</i> , but only related to <i>anxiety-withdrawal in girls</i>

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Kaplan et al., (1997)	198 abused adolescents (12–18 yrs); 99 physically abused; 99 nonabused (50%F; 100% W)	C/S	physical abuse (state records), Parental Bonding Instrument; FACES; # parents in home, separation from parent, head trauma	depressive disorders, disruptive disorders, substance abuse (SADS for children, Connors' Teacher Rating Scale) PTSD (SKID-NP), available school & medical info (int & ext)	<b>Mixed Specificity for physical abuse:</b> for current diagnoses, abuse increased the risk for being diagnosed with <i>unipolar depressive disorders, disruptive disorders, &amp; cigarette smoking</i> , but not ADHD or PTSD; For lifetime disorders, abuse increased risk of major depression, dysthymia, conduct disorder, drug abuse or dependence, & cigarette smoking, but not ADHD or PTSD
Kolko, Moser, & Weldy (1988)	103 inpatient children (5–14 yrs); 29 sexually abused, 52 physically abused (27% F; 72% W, 28% AA)	C/S	sexual abuse, physical abuse (medical chart record), physical exams & Sexual Abuse Symptom Checklist (dev au)	parent report sx of sexual activity, anxiety, depression, conduct problems, withdrawal (SASC) & aggression, sexual behavior, anxiety, withdrawal, sleep problems, depression (hospital chart review) (int & ext)	<b>Specificity for sexual abuse int (not ext) &amp; sexual acting out:</b> sexual abuse was specifically related to <i>sexual activity, anxiety, withdrawal, depression</i> <b>No Specificity for physical abuse</b>
Livingston, Lawson, & Jones (1993)	41 clinic-referred children (6–15 yrs); 26 sexually abused, 15 physically abused (44% F; 52% AA, 48% W)	C/S	sexual abuse, physical abuse (clinic reports of abuse)	self-report behavior disorders, somatization symptoms, PTSD, depressive disorders, anxiety disorders, psychotic symptoms (DICA-6-R) (int & ext)	<b>Mixed Specificity for sexual abuse:</b> sexual abuse associated with ADHD, <i>separation anxiety disorder, PTSD, somatization, and one psychotic symptom</i> ; only one psychotic symptom was found specifically related to sexual abuse
Murata (1994)	23 children (8–12 yrs; 100%M; 91% AA, 9% W)	C/S	mother's use of violence (CTS), family stresses (FILE)	parent and teacher report internalizing and externalizing syndromes (TRF, CBCL) (int & ext)	<b>No Specificity for physical abuse</b> <b>Specificity for int (not ext) for physical abuse:</b> mother's use of violence & family stress were associated with internalizing problems (but not externalizing)
Prino & Peyrot (1994)	68 children (5–8 yrs): 21 physically abused (52% F; 57% AA, 43% W), 26 neglected (27%F; 58% W, 42% AA), & 21 controls (52%F; 57% AA, 43%W)	C/S	physical abuse, neglect (medical records, child protective services, court documented)	teacher reports aggressive and withdrawn behavior (PASS) (int & ext)	<b>Specificity for ext (not int) for physical abuse:</b> physical abuse was specifically related to <i>aggressive behavior</i> (compared to neglect or control); <b>Specificity for int (not ext) for neglect:</b> neglect was specifically related to <i>withdrawn behavior</i> (compared to abuse and control)

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Rivinus, Levoy, Matzko, & Seifer (1992)	109 children (4–14 yrs; 23% F; NFDP) from psychiatric inpatient unit	C/S	parental substance abuse, physical &/or sexual abuse (intake, clinical information)	clinician report of DSM-III-R diagnoses (clinical evaluation) (int & ext)	<b>Mixed Specificity for physical and sexual abuse:</b> <i>physical &amp;/or sexual abuse</i> was associated with <i>dysphoric mood, oppositional symptoms, PTSD &amp; Borderline Personality Disorder, children of parental substance abusers &amp; abused</i> were more likely to be diagnosed with <i>schizophrenia or ODD; parental substance abuse</i> was associated with <i>enuresis, obsessive-compulsive symptoms &amp; learning disabilities</i>
Sadeh, Hayden et al. (1994)	100 inpatient children (2–13 yrs; 17% F; NFDP)	C/S	sexual abuse &/or physical abuse (from child & parent interview)	self-reported depression symptoms (CDI), DSM-III-R diagnoses (interview) (int & ext)	<b>Specificity for sexual abuse int (not ext);</b> <i>sexual abuse</i> was specifically related to <i>PTSD</i> (compared to physical abuse or no abuse) & <i>sleep problems</i> (i.e., parasomnias)(not depression) <b>No Specificity for physical abuse</b>
Sibthorpe, Drinkwater et al. (1995)	155 Australian youth (12–17 yrs; 46% F; 7% Aborigine, 93% O)	C/S	self-report sexual & physical abuse interview questions (dev by au)	self-report substance use & suicidality sx (interview dev by au) (int & ext)	<b>Specificity for sexual abuse int (not ext)</b> sexually (but not physically abused) participants were more likely, than those not abused, to have attempted suicide <b>No Specificity for physical abuse</b>
Silverman, Reinherz, & Giaconia (1996)	375 adolescents (15 yrs at time 1 & 21 at time 2; 50% F; 99% W, 1% O)	L (2 ts, 6 yrs apart) & C/S	physical abuse, sexual abuse (clinical interview)	self-reported psychological syndromes (YSR), depressive symptoms (CDI) anxiety (PHSC-S) & suicidal ideation (CDI) (int & ext)	<b>Mixed Specificity for physical and sexual abuse:</b> <i>sexual abuse</i> was associated with <i>somatic complaints, anxiety-depression, social problems, thought problems, attention problems, aggressive behavior, depressive symptoms, &amp; suicidal ideation; physical abuse</i> was associated with suicidal ideation (boys), & withdrawal, somatic complaints, anxiety-depression, thought problems, attention problems, delinquent behavior, aggressive behavior (YSR) & symptoms of depression, anxiety, & suicidal ideation (girls)
Threlkeld & Thyer (1992)	117 children in tx (<17 yrs; 46% F; 55% W, 38% AA, 7% O)	C/S	sexual abuse & physical abuse (record review)	dSM diagnoses (no method identified) (int & ext)	<b>No Specificity for physical or sexual abuse</b>
White, Halpin, Strom, & Santilli (1988)	58 children (26 yrs): 17 sexually abused, 18 neglected, 23 controls (NFDP)	C/S	sexual abuse, neglect (hospital-referred)	sexual acting out, somatic complaints, withdrawal, oppositional (MCIDI) (int & ext)	<b>Specificity for sexual acting out:</b> <i>sexual abuse</i> was specifically associated with <i>sexual acting out</i> (but not other symptoms) <b>No Specificity for neglect</b>

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Williamson, Borduin, & Howe (1991)	50 adolescents (12-17 yrs; 82% F); 12 neglected, 12 physically abused, 15 sexually abused, 11 control (75-92% W, 8-25% AA)	C/S	neglect, physical abuse, sexual abuse (Juvenile Court)	parent report of total sx (SCL-90-R), conduct, aggression, anxiety-withdrawal and attention problems (RBPC) (int & ext)	<p><b>Specificity for ext (not int) for physical abuse:</b> physical abuse was specifically associated with conduct problems and socialized aggression (compared to control); all 3 types of abuse (compared to control) were associated with attention problems</p> <p><b>Mixed Specificity for sexual abuse:</b> sexual abuse was associated with conduct problems and anxiety-withdrawal (compared to control)</p> <p><b>Specificity for ext (not int) for neglect:</b> Neglect was associated with conduct problems (compare to control) and socialized aggression (compared to sexual abuse and control)</p>
See also Lynch & Cicchetti (1998); McCloskey, Figueredo, & Koss (1995); O'Keefe (1994, 1997); Sternberg et al., (1993) & Tang (1997) in Exposure to Violence					
<b>Divorce/Marital Conflict</b>					
David, Forchard, & Armistead (1996)	146 adolescents (11-15 yrs; 52% F; NFDP)	L (2 ts, 1 year apart)	family conflict, (FES); witness of marital conflict (OPS), marital dissatisfaction (DAS)	parent & teacher report ext & int sx (RBPC) (int & ext)	<p><b>Mixed Specificity:</b> family conflict more associated with child problems than marital satisfaction; family conflict &amp; witnessed marital conflict (but not marital satisfaction) predicted externalizing problems at time 1 &amp; 2 &amp; internalizing problems at time 1; witnessed marital conflict only predicted internalizing at time 2 (mother report); marital satisfaction predicted externalizing problems at time 2; witnessed marital conflict predicted internalizing problems at time 1 (teacher report)</p>
Jenkins & Smith (1990)	119 British children (9-12 yrs); poor marriages (57) & good marriages (62)	C/S	overt parental conflict, covert tension, discrepancy in reading (MMRS interview)	31 int & ext behavioral sx, parent report & child report (interview dev by Graham & Rutter plus ICD9 criteria)(int & ext)	<p><b>Specificity for ext (not int):</b> there were more correlations between overt parental conflict &amp; child behavior problems than for covert tension or discrepancy in child rearing practices, child reports indicated that all types of conflict were related to externalizing problems, but not internalizing problems (no differences for parent reports)</p>

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Jouriles, Norwood, & Vincent (1996a)	55 children (5–12 yrs; 66% F; 9.5% W, 5% O) from families seeking marital therapy	C/S	marital aggression (physical, verbal, & behavioral) (CTS)	parent-report externalizing and internalizing behavior (conduct & personality subscales of the BPC) (int & ext)	<b>Specificity for ext (not int):</b> physical marital aggression and other forms of marital aggression both related to externalizing (but not to internalizing problems); stressor specific: other forms of aggression were more specifically related to externalizing problems than physical aggression
Jouriles, Norwood, & Vincent (1996b)	199 children (5–12 yrs; 47% F; 39% W, 24% AA, 37% L) in battered women's shelters	C/S	marital aggression, divided into physical violence & other forms of aggression (CTS)	parent report (CBCL) (int & ext)	<b>Mixed Specificity:</b> physical aggression was specifically related to externalizing & other forms of marital aggression were specifically related to internalizing
Marttunen, Aro, Henriksson, & Lonqvist (1994)	53 adolescents (13–19 yrs; 17% F; 100% Finnish) who committed suicide	C/S	parental violence, separations, conflicts, parental divorce, (archival records, parent-report stress interviews, clinician-report stress interviews) (dev by au)	composite of archival, clinician & parent-report depressive disorder, substance abuse & other disorders (int & ext)	<b>Specificity for ext (not int):</b> parental violence was more common among alcohol abusing victims than among depressive victims; <i>interpersonal separations</i> were more common among alcohol abusing victims than depressive victims; <i>Specificity by gender:</i> interpersonal conflicts were more common among alcohol-abusing male (but not female) victims than depressive victims
See also Conger et al. (1992, 1993, 1994) in Poverty					
<b>Poverty</b>					
Conger, Conger, Elder, Jr., Lorenz, Simons, & Whitbeck (1992)	205 adolescents (7 <sup>th</sup> grade) (100% M; 100% W)	C/S	parent-report economic hardship scale, economic pressure; observer-report marital conflict measure (dev by au)	self-report hostility symptoms (modified BDHS), int sx (SCL-90 depression & hostility subscales); sibling-report hostility sx (dev by au) (int & ext)	<b>Mixed Specificity for poverty:</b> economic hardship indicator 'debts to assets ratio' was associated with sibling-report hostility symptoms, economic pressure indicators 'make ends meet', 'material needs' & 'economic adjustments' were associated with self-report internalizing symptoms & economic pressure indicator 'material needs' was associated with sibling-report hostility symptoms <b>Specificity for ext (not int) for marital conflict:</b> marital conflict was associated with self-report (but not sibling-report) hostile symptoms (but not self-report internalizing symptoms)

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Conger, Conger, Elder, Jr., Lorenz, Simons et al. (1993)	220 adolescents (12–14 yrs, 7 <sup>th</sup> grade; 100% F; 100% W)	C/S	parent-report economic hardship scale, economic pressure; observer-report marital conflict (dev by au)	self-report hostility sx (modified BDHS), int sx (SCL-90 depression, hostility, anxiety, somatization subscales); sibling-report hostility sx (dev by au) (int & ext)	<b>Specificity for ext (not int) for poverty:</b> economic hardship indicator 'debts to assets ratio' was associated with self-report hostility symptoms, economic hardship indicator 'income loss' was associated with sibling-report hostility symptoms; <b>No Specificity for marital conflict:</b> marital conflict was associated with all three symptom measures
Conger, Ge, Elder, Jr., Lorenz, & Simons (1994)	378 adolescents (7 <sup>th</sup> grade; 52% F; 100% W)	L (3 ts, 1 year apart)	parent-report economic hardship scale, economic pressure; marital conflict; observer-report marital conflict (dev by au)	self-report depression & anxiety sx (SCL-90 & NEO-personality anxiety & depression subscales), ext sx (modified BDHS, SCL-90 & NEO-personality hostility subscales), & aggression (ABS-dev by au) (int & ext)	<b>No Specificity for poverty:</b> no poverty indicators were associated with symptoms; <b>Mixed Specificity for marital conflict:</b> father-report marital conflict was associated with trait hostility (ext), & trait anxiety (int); observer-report marital conflict was associated with 3 of the 4 externalizing measures (all but self-report aggressive behavior)
See also Levendosky & Graham-Bermann (1998) in Exposure to Violence					
<b>Illness</b>					
Wallander, Varni, Babani, Banis, & Wilcox (1988)	270 children: 80 diabetes, 77 spina bifida, 40 hemophilia, 30 chronic obesity, 24 juvenile rheumatoid arthritis, & 19 with cerebral palsy	C/S	established illness (patients recruited based on physician diagnosis)	parent-report int & ext sx (CBCL) (int & ext)	<b>Specificity:</b> no differences between groups on internalizing; juvenile rheumatoid arthritis lower on externalizing than other groups; cerebral palsy group lower on social competence
<b>Cumulative Stress</b>					
Aseltine, Jr., & Gore (1993)	918 adolescents (9 <sup>th</sup> –11 <sup>th</sup> grade; 59% F; 100% W)	L (2 ts, 1 year apart)	self-report family relationship stress scale, friend relationship stress scale, graduation status (dev by au)	self-report depressive sx (CES-D), alcohol use (dev by au), delinquency (modified BDS) (int & ext)	<b>Specificity: family &amp; friend stress</b> predicted depressive symptoms (but not alcohol abuse symptoms); <i>friend</i> (but not family) stress predicted increases in <i>delinquent</i> symptoms; <i>graduation</i> from high school predicted increases in <i>alcohol</i> abuse symptoms, but <i>decreases</i> in depressive symptoms

Table 1 Continued

Author	Sample	Design	Stress measure	Outcome measure	Findings
Lewinsohn, Gotlib, & Seeley (1995)	1507 adolescents (14–18 yrs at Time 1; 54% F; 91% W, 9% O)	L (2 ts, 1 year apart)	self-report conflict with parents & parental death (psychosocial interview dev by au)	self-report major depressive disorder & substance abuse disorder (K-SADS) (int & ext)	<b>Mixed Specificity:</b> those with <i>higher rates of conflict</i> with parents (but not parental death) were more likely to be <i>depressed</i> & <i>use substances</i>
Sandler, Reynolds, Kliewer, & Ramirez (1992)	359 youth (8–16 yrs; 43% F; NFD), 92 who had lost a parent, 94 whose parents had divorced, 99 w/ asthma, 74 controls	C/S & L	self-report cumulative stress scale (GLESC) & interview (dev by au); events categorized as 'conflict events' & 'loss events'	self-report depression (CDI) & conduct disorder sx (YHS) (int & ext)	<b>Specificity:</b> conflict (but not loss) events were associated with conduct (but not depression) sx; loss (but not conflict) events were associated with depression (but not conduct) sx; opposite pattern in sample of youth whose parents had recently divorced

See also Attar, Guerra, & Tolan (1994) in Exposure to Violence

#### General Abbreviations Key

**F** = female, **M** = male; **NFDP** = no further demographic information provided; **ts** = Data Collection Time Points; **C/S** = Cross-sectional; **L** = Longitudinal; **W** = White/European American, **AA** = African American, **L** = Latino, **AS** = Asian-American, **AI** = American Indian, **O** = Other.

#### Stress Measures Key

**CTS** = Conflict Tactics Scale; **CVS** = Community Violence Survey; **DAS** = Dyadic Adjustment Scale; **dev by au** = developed by authors; **FES** = Family Environment Scale; **FILE** = Family Inventory of Life Events; **GLESC** = General Life Events Schedule for Children; **HCSI** = Hispanic Children's Stress Inventory; **HURTE** = Hurricane-Related Traumatic Experiences; **LEQ** = Life Events Questionnaire; **LER** = Life Events Record; **MMRS** = Maudsley Marital Rating Scale; **OPS** = O'Leary Porter Scale; **PYS** = Pittsburgh Youth Study; **SCSS** = Survey of Children's Stress Symptoms; **SECV** = Survey of Exposure to Community Violence; **SI** = Stress Index; **VAWS** = Violence Against Women Scale.

#### Outcome Measures Key

**ABS** = Aggressive Behavior Scale; **BDHS** = Buss & Durkee Hostility Scale; **BDS** = Bachman Delinquency Scale; **BORRTI** = Bell Object Relations & Reality Testing Inventory; **BPI** = Behavior Problems Index; **CAS** = Child Assessment Schedule; **CBCL** = Child Behavior Checklist; **CBCL-PTSD** = Child Behavior Checklist Post-traumatic Stress Disorder; **CCDS** = Checklist of Child Distress; **CDC** = Child Dissociative Checklist; **CDI** = Child Depression Inventory; **C-SBI** = Child Sexual Behavior Inventory; **DIS** = Diagnostic Interview Schedule; **DICA** = Diagnostic Interview for Children & Adolescents; **DICA-R** = Diagnostic Interview for Children & Adolescents Revised; **DSRS** = Depression Self-Rating Scale; **DYSSRDQ** = Denver Youth Study Self Reported Delinquency Questionnaire; **ext** = externalizing; **GI** = Grief Index; **int** = internalizing; **JMAS** = Jesness Manifest Aggression Scale; **K-SADS** = Schedule for Affective Disorders and Schizophrenia; **MCDI** = Minnesota Child Development Inventory; **MMRS** = Maudsley Marital Rating Scale; **PASS** = Pittsburgh Adjustment Survey; **PBQ** = Preschool Behavior Questionnaire; **PHSC-S** = Piers-Harris Self Concept Scale; **P-PTSD-S** = Purdue PTSD Scale; **PTSD-RI** = PTSD Reaction Index; **PTSD-SS** = PTSD Symptom Scale; **R** = Rorschach; **RBPC** = Revised Behavior Problem Checklist; **RCMAS** = Revised Children's Manifest Anxiety Scale; **SASC** = Sexual Abuse Symptom Checklist; **SCL** = Symptom Checklist; **STAIC-1F** = State Trait Anxiety Inventory for Children-Trait Form; **TRF** = Teacher Report Form; **TSC-C** = Trauma Symptom Checklist-Children; **WVPS** = Weiss Peters Scale; **YHS** = Youth Hostility Scale; **YSR** = Youth Self Report.

## Exposure to violence

Sixteen Stressor-Outcome specific studies tested for specific relations between exposure to violence and particular internalizing and externalizing psychological outcomes (see Table 1). Exposure to the following specific types of violence has been examined: domestic violence, community violence, and war violence. Outcomes examined include PTSD, depression, general anxiety, and aggression symptoms, and broad-band internalizing and externalizing syndromes. Unfortunately, few of these studies used the same measures of stressors and outcomes, thus limiting conclusions that may be drawn.

To examine general patterns across this disparate body of studies, we grouped outcomes into internalizing (i.e., anxiety, depression, somatization, withdrawal, PTSD) and externalizing (i.e., aggression, delinquency, attention problems, oppositionality, substance abuse) categories. Of the sixteen *Stressor-Outcome specific* studies examining both internalizing and externalizing outcomes, one-half of the studies reported evidence of mixed specificity or no specificity for particular outcomes (Gorman-Smith & Tolan, 1998; Jouriles, Barling, & O'Leary, 1987; Jouriles, Norwood, McDonald, Vincent, & Mahoney, 1996, study 2; Lynch & Cicchetti, 1998; McCloskey, Figueiredo, & Koss, 1995; O'Keefe, 1994a, 1997; Tang, 1997). Five of the eight reporting specificity reported a specific relation between exposure to violence and *externalizing* outcome (Attar, Guerra, & Tolan, 1994; Durant, Pendergrast, & Cadenhead, 1994; Jouriles et al., 1996, study 1; Marttunen, Aro, Henriksson, & Lonnqvist, 1994; Sternberg et al., 1993), and three reported a specific relation between exposure to violence and internalizing outcome (Levendosky & Graham-Bermann, 1998; Osofsky, Wewers, Hann, & Fick, 1993; Rossman, Bingham, & Emde, 1997). Methodological explanations for discrepancies across studies are not readily apparent. For example, the four studies using the most similar methodology (all four used a version of the Conflict Tactic Scale and CBCL and examined a sample of predominantly pre-adolescent children) reported inconsistent results.

Most *Outcome specific* studies of exposure to violence failed to find evidence of specificity (Brent, Perper, Moritz, Allman et al., 1993; O'Keefe, 1994b; Pfeffer et al., 1997; Spaccarelli, Sandler, & Roosa, 1994; Brent, Moritz, Bridge, Perper, & Canobbio, 1996; Schwab-Stone et al., 1995). However, two outcome-specific studies reporting specificity effects found specificity for *internalizing* (Brent, Perper, Moritz, Friend et al., 1993; Freeman, Shaffer & Smith, 1996) while one found specificity for *externalizing* outcomes (Cooley-Quille, Turner, & Beidel, 1995), which suggests a pattern inconsistent with the one found in Stressor-Outcome specific studies. A theory-based explanation for these discrepant findings is suggested by the fact that both of the studies reporting specificity for internalizing

symptoms focused on experiences that may also be conceptualized as loss events (i.e., *murder* of a sibling; Freeman et al., 1996, and *suicide* of a peer; Brent, Perper, Moritz, Friend et al., 1993).

## Abuse

Thirty-three Stressor-Outcome specific studies tested for specific relations between child abuse and particular internalizing and externalizing psychological outcomes (see Table 1). Specific types of abuse examined include physical abuse, sexual abuse, and neglect. In most cases, these particular types of abuse were compared with one another in relation to several specific outcomes. Abuse, as a whole, was rarely examined in relation to other categories of stress. Specific outcomes examined include suicidality, substance abuse, sexual acting out, delinquency, aggression, attention problems, general anxiety, PTSD, withdrawal, somatization, eating disorders, depression symptoms, and broad-band internalizing and externalizing syndromes. As with exposure to violence studies, few of these studies used the same measures of stressors and outcomes, thus limiting conclusions that may be drawn.

Of the 33 *Stressor-Outcome specific* studies focusing on abuse, 31 studies examined internalizing and externalizing outcomes related to *physical abuse*. Of these studies, seven studies demonstrated evidence for specificity, with four reporting specificity for *externalizing* outcomes (Ackerman, Newton, McPherson, Jones, & Dykman, 1998; Feldman et al., 1995; Prino & Peyrot, 1994; Williamson, Borduin, & Howe, 1991), and three reporting specificity for internalizing outcomes (Crittenden, Claussen, & Sugerman, 1994; DePaul & Arruabarrena, 1995; Murata, 1994). Results of *Outcome specific* studies reveal a similar pattern of findings. Four of these studies, which examined both internalizing and externalizing outcomes, failed to find evidence of specificity (Glod & Teicher, 1996; Flisher, Kramer, Hoven, & Greenwald, 1997; Trickett, 1993; Pelcovitz et al., 1994). The two studies that found evidence of specificity reported evidence of specificity for *externalizing* outcomes (Hennessy, Rabideau, Cicchetti, & Cummings, 1994; Walker, Downey & Bergman, 1989).

Nineteen *Stressor-Outcome specific* studies examined both internalizing and externalizing outcomes in relation to *sexual abuse*. Twelve of these failed to find evidence of specificity (Ackerman et al., 1998; Bayatpour, Wells, & Holford, 1992; Boney-McCoy & Finkelhor, 1995; Caviola & Schiff, 1988; Cohen et al., 1996; Hernandez, 1995; Livingston, Lawson, & Jones, 1993; Rivinus, Levoy, Matzko, & Seifer, 1992; Silverman, Reinhertz, & Giaconia, 1996; Threlkeld & Thyer, 1992; White, Halpin, Strom, & Santilli, 1988; Williamson et al., 1991). Among those reporting specificity, six reported specificity for *internalizing* outcomes (Dykman et al.,



1997; Friedrich, Jaworski, Huxsahl, & Bengston, 1997; Gale, Thompson, Moran, & Sack, 1988; Kolko, Moser, & Weldy, 1988; Sadeh, Hayden, McGuire, Sachs, & Civita, 1994; Sibthorpe, Drinkwater, Gardner, & Bammer, 1995), and one (which included only males) reported specificity for externalizing outcomes (Hernandez, Lodico, & DiClemente, 1993). Although these studies suggest a pattern of increased risk for internalizing outcomes, girls are at substantially higher risk both for sexual abuse and for internalizing symptoms (Berliner & Elliott, 1996). Thus, the association between sexual abuse and internalizing outcomes may simply reflect this well-established relation. Unfortunately, the vast majority of studies reviewed above were not designed to maximally examine specificity hypotheses. Most of these studies failed to take gender into account in their examination of specificity effects (i.e., studies reporting heightened rates of internalizing symptoms among sexually abused, relative to physically abused, youth failed to control for differences in gender representation across the two samples).

Results of *Outcome specific* studies of sexual abuse provide little evidence of specificity. Thirty of 35 studies that examined both internalizing and externalizing outcomes failed to find evidence of specificity (Black, Dubowitz, & Harrington, 1994; Brand, King, Olson, Ghaziuddin, & Naylor, 1996; Chandy, Blum, & Resnick, 1996; Cohen & Mannarino, 1988; Cosentino, Meyer-Bahlburg, Alpert, Weinberg, & Gains, 1995; Einbender & Friedrich, 1989; Garnefski & Arends, 1998; Glod & Teicher, 1996; Goldston, Turnquist, & Knutson, 1989; Hazzard, Celano, Gould, Lawry, & Webb, 1995; Hibbard & Hartman, 1992; Inderbitzen-Pisaruk, Shawchuck, & Hoier, 1992; Kuhn, Arrelano, & Chavez, 1998; Kumar, Steer, & Deblinger, 1996; Luster & Small, 1997; Mannarino, 1988; Mannarino & Cohen, 1996; Martin, 1996; McClellan, Adams, Douglas, McCurry, & Storck, 1995; McLeer, Deblinger, Henry, & Orvaschel, 1992; McLeer, Callaghan, Henry, & Wallen, 1994; Shapiro, Leifer, Marone, & Kassem, 1992; Smith & Howard, 1994; Stern, Lynch, Oates, O'Toole, & Cooney, 1995; Tong, Oates, & McDowell, 1987; Watts & Ellis, 1993; Wells, McCann, Adams, Voris, & Dahl, 1997; Wolfe, Gentile, & Wolfe, 1989; Young, Bergandi, & Titus, 1994). The few which reported specificity effects split slightly in favor of *internalizing* (Johnson & Kenkel, 1991; Mian, Marton, & LeBaron, 1996; Hussey, Strom, & Singer, 1992) over externalizing outcomes (Hussey & Singer, 1993; Rotheram-Borus, Mahler, Koopman, & Langabeer, 1996).

Due to the significant attention paid to the relation between sexual abuse and PTSD, we examined studies testing for specificity between sexual abuse and this particular psychological outcome. The majority of studies conducted in this area suggest there is a specific relation between sexual abuse and PTSD. All eight of the Stressor-Outcome studies (100%) found specificity in the relation between sexual abuse and

PTSD (Deblinger, McLeer, Atkins, Ralph, & Foa, 1989; Dykman et al., 1997; Friedrich et al., 1997; Haviland, Sonne, & Woods, 1995; Livingston et al., 1993; Rivinus et al., 1992; Sadeh et al., 1994; Timmons-Mitchell, Holtz, & Semple, 1997). In addition, of the three Outcome specific studies that examined the relation between sexual abuse and PTSD, two found evidence of specificity (McClellan et al., 1995; McLeer et al., 1992) while one study did not find a specific relation between sexual abuse and PTSD (Berliner & Conte, 1995). Although the findings are quite consistent, methodological concerns related to gender-specific effects, highlighted above, temper conclusions that may be drawn.

Several sexual abuse studies also tested the hypothesis that sexual abuse is specifically associated with 'sexual acting out'. As this outcome has varied from conceptualization as an internalizing outcome (symptom of PTSD; Deblinger et al., 1989) to conceptualization as an externalizing outcome (an aggressive act; Dykman et al., 1997), we chose to examine it separately from these broad-band classification systems. Eight *Stressor-Outcome specific* studies tested for specificity for *sexual acting out*. Six (75%) found evidence of *specificity* (Deblinger et al., 1989; Friedrich et al., 1997; Gale et al., 1988; Higgins & McCabe, 1998; Kolko et al., 1988; White et al., 1988) and two did not (Dykman et al., 1997; Hernandez et al., 1993). Fifteen *Outcome specific* studies tested for specificity for sexual acting out. Twelve of these (80%) found a *specific relation* between sexual abuse and sexual acting out behaviors (Cohen & Mannarino, 1988; Cosentino et al., 1995; Einbender & Friedrich, 1989; Hibbard & Hartman, 1992; Inderbitzen-Pisaruk et al., 1992; Mannarino, 1988; Mannarino & Cohen, 1996; Mian et al., 1996; Rotheram-Borus et al., 1996; Smith & Howard, 1994; Wells et al., 1997; Young et al., 1994), and three failed to find evidence of specificity for sexual abuse and sexual acting out (Berliner & Conte, 1995; Chandy et al., 1996; Stern et al., 1995).

A small handful of *Stressor-Outcome specific* studies examined both internalizing and externalizing outcomes in relation to *neglect*. Two failed to find evidence of specificity (Crittenden et al., 1994; White et al., 1988). Of those reporting specificity effects, two reported specificity for *externalizing* outcomes (DePaul & Arruabarrena, 1995; Williamson et al., 1991), and one reported specificity for internalizing outcomes (Prino & Peyrot, 1994). The two Outcome specific studies to examine both internalizing and externalizing outcomes failed to find evidence of specificity (Famularo, Kinscherff, & Fenton, 1992; Markward, 1997).

### Divorce/Marital conflict

Eight Stressor-Outcome specific studies tested for specific relations between divorce (or marital

conflict) and particular internalizing and externalizing psychological outcomes (see Table 1). Most of these studies have focused on marital conflict, rather than divorce, *per se*. Specific measures of marital conflict have included parent-report measures of inter-parental conflict, child-report measures of witnessing conflict, and observer-report ratings of marital interactions. Specific outcomes examined include depression, substance abuse, anxiety, somatization, aggression, hostility, antisocial, and hyperactivity symptoms, and broad-band internalizing and externalizing syndromes. As with the exposure to violence and child abuse studies, few of these studies used the same measures of stressors and outcomes, thus limiting conclusions that may be drawn.

One-half (four of eight) of the divorce/marital conflict studies reported evidence of specificity for *externalizing* outcomes (Conger et al., 1992; Jenkins & Smith, 1991; Jouriles et al., 1996; Marttunen et al., 1994). As with most of the specificity studies reviewed, variability in methodology hampers comparisons across studies for the purpose of understanding discrepancies in outcomes. However, the series of Conger studies, using similar methodologies in differing samples (Conger et al., 1992, 1993, 1994), provides an exception to this pattern. Conger and colleagues found evidence of specificity between observer-report marital conflict and externalizing outcomes in their all-male sample (Conger et al., 1992) and in their mixed sample (52% female; Conger et al., 1994), but not in their all-female sample (Conger et al., 1993). This pattern fits with recent findings that girls exhibiting heightened rates of externalizing symptoms are particularly at risk, relative to male counterparts, for co-occurring internalizing symptoms (O’Koon, 1999). Although the Conger studies do not define themselves as specificity studies, the incremental, programmatic nature of their research provides an excellent example of the type of research needed in this area.

*Outcome specific* studies on divorce/marital conflict provide an interesting contradiction. Although most (six) of those examining both internalizing and externalizing outcomes failed to find specificity effects (Fergusson, Horwood, & Lynskey, 1994; Forehand, Neighbors, & Wierson, 1991; Hoyt, Cowen, Pedro-Carroll, & Alpert-Gillis, 1990; Rodgers, 1994; Reese & Roosa, 1991; West, Sandler, Pillow, Baca, & Gersten, 1991), those reporting specific associations were split, slightly, in favor of *internalizing* (Brody & Forehand, 1990; Forehand, McCombs, Long, Brody, & Fauber, 1988; Long, Slater, Forehand, & Fauber, 1988) over externalizing outcomes (Gould, Shaffer, Fisher, & Garfinkel, 1998; Smith, Howard, & Monroe, 1998).

Although these findings appear contradictory, examination of the studies reporting specificity for internalizing outcomes provides evidence of an interesting pattern. All found evidence of specificity between marital conflict and internalizing symptoms

within samples of youth who had *recently* experienced their parents’ divorce. This pattern is consistent with theory-based work by Sandler and colleagues (reviewed below), which suggests that marital conflict, in the context of recent divorce, may be associated, in meaning, with separation and loss events and, thus, ‘pull for’ internalizing outcomes (Sandler et al., 1992).

### Stressors from various categories

Remaining Stressor-Outcome specific studies are spread thinly across several categories (poverty, illness, cumulative stress). Four *Stressor-Outcome specific* studies have included *poverty* as a stressor, examining both internalizing and externalizing outcomes (see Table 1). Among these studies, three (Conger et al., 1992, 1994; Levendosky et al., 1998) failed to find evidence of specificity, and one (Conger et al., 1993) found evidence of specificity for *externalizing* outcomes. Most *Outcome specific* poverty studies also failed to find specific effects (Bolger, Patterson, Thompson, & Kupersmidt, 1995; DuBois, Felner, Meares, & Krier, 1994; Elder, Jr., Conger, Foster, & Ardel, 1992; Felner et al., 1995; Kolko & Kazdin, 1993; McLoyd, Jayaratne, Ceballo, & Borquez, 1994; Mezzich et al., 1997; Reinherz et al., 1993). Those that reported specificity findings found evidence of specificity for *internalizing* outcomes (Brody & Flor, 1997; Masten, Miliotis, Graham-Bermann, Ramirez, & Neemann, 1993; Lempers, Clark-Lempers, & Simons, 1989).

Only one Stressor-Outcome illness study has been conducted (see Table 1), and most Outcome specific studies in this area, which examined both internalizing and externalizing outcomes, failed to find evidence of specificity (Burke et al., 1989; Engstrom, 1992; Kashani, Konig, Sheppard, Wilfley, & Morris, 1988; Mikail & Von Baeyer, 1990; Siegel et al., 1992). Those reporting evidence of specific effects were evenly split between specificity for internalizing (Garralda, Jameson, Reynolds, & Postlethwaite, 1988) and externalizing outcomes (Smith et al., 1998).

There were no Stressor-Outcome specific disaster (i.e., natural or human-made disaster) studies, and most *Outcome specific* disaster studies, which examined internalizing and externalizing outcomes, failed to find evidence of specificity (Durkin, Khan, Davidson, Zaman, & Stein, 1993; Laor et al., 1996; La Greca, Silverman, & Wasserstein, 1998; March, Amayaj-Jackson, Terry, & Costanzo, 1997; Shaw et al., 1995). All three of those which did find evidence of specificity, reported specificity for *internalizing* outcomes (Breton, Valla, & Lambert, 1993; Johnston, Gonzalez, & Campbell, 1987; Najarian, Goenjian, Pelcovitz, Mandel, & Najarian, 1996).

None of the Stressor-Outcome specific cumulative stress studies examined similar categories of stress

(see Table 1), precluding comparisons across studies. Nonetheless, one Stressor-Outcome specific cumulative stress study merits discussion, as it provides a prototype of theory-based specificity research. Sandler et al. (1992) tested two theory-based specificity hypotheses: (1) stressful events involving separation from important family members will be specifically associated with internalizing symptoms, and (2) stressful events involving conflict will be specifically associated with externalizing symptoms. These hypotheses were tested in four samples of children (a group of children who had experienced the death of a parent, a group who had experienced their parents' divorce, a group of children with chronic asthma, and a control group of children who had not experienced any of these stressors) to examine the interaction of specificity findings with varying environmental contexts.

Sandler and colleagues (1992) reported partial support for their hypotheses. Conflict events were associated with externalizing symptoms, but not internalizing symptoms, in two of their samples (parental death and control samples), and separation events were associated with internalizing symptoms, but not externalizing symptoms, in two of their samples (parental death and asthma samples). However, findings directly contradictory to the authors' hypotheses were found among the group of children who experienced divorce. In this group, conflict events were associated with internalizing and separation events were associated with externalizing symptoms. These findings illustrate the importance of integrating specificity and moderator designs, as the differing contexts of the various samples in this study moderated the specific relations between particular stressors and particular outcomes. Sandler and colleagues (1992) propose that the differences in specificity findings across different samples may reflect differences in meaning applied to the stressful events for different groups of children. In particular, children of divorced parents may respond differently from other children to conflict and separation events because of the inter-relatedness of these two experiences within the context of divorce. This hypothesis is consistent with the findings, reported in the divorce/marital conflict section, that marital conflict was associated with internalizing symptoms among youth recently exposed to their parents' divorce (Forehand et al., 1988; Forehand et al., 1988; Long et al., 1988).

Related to Sandler and colleagues' study (1992) is an Outcome specific study, which represents the sole study to include specificity of moderator in their theory-based specificity design. Shirk, Boergers, Eason, and Van Horn (1998) hypothesized that interpersonal stress would predict depression, but not hostility, symptoms in a sample of 8th grade youth and that interpersonal schema would moderate the relation between interpersonal stress and depression, but not hostility, symptoms. The authors report

support for both their hypotheses. Interpersonal stress was specifically predictive of depressive symptoms, and interpersonal schema moderated the relation between interpersonal stress and depression (but not hostility) symptoms.

### Summary and integration of findings

Results of studies testing the hypothesis that particular stressors are specifically related to internalizing or externalizing outcomes provide little support for this hypothesis. In general, the results of this review are more consistent with tenets of equifinality and multifinality in the relation between stress and psychological problems in children and adolescents than they are of specificity. Across the various stressors examined (exposure to violence, abuse, divorce/marital conflict, poverty, illness, and cumulative stress), the most consistent evidence for specificity was found in relation to sexual abuse. Sexual abuse was associated specifically with internalizing outcomes, PTSD, and with sexual acting out across several studies. Interestingly, the outcomes of PTSD and sexual acting out are the two areas in this review that are more specific in nature, and these are the outcomes in which there was evidence of specificity. These findings may reflect that researchers examining sexual abuse are interested in these particular theoretically based outcomes (PTSD and sexual acting out), and because of this interest, there were enough comparable studies examining these specific relationships to reveal patterns of specificity.

The pattern of findings demonstrating a specific relation between sexual abuse and internalizing outcomes should be interpreted with caution, given the fact that this association may reflect heightened risk for both sexual abuse and internalizing outcomes among girls. Dominant theoretical models of sexual abuse (Finkelhor & Brown, 1985; Spaccarelli, 1994) argue that sexual abuse is linked with both internalizing and externalizing outcomes, through varying mediating processes (e.g., emotions, cognitions, coping), and that these relations are moderated by particular environmental contexts (e.g., offender-victim relationships, family relationships/responses and community contexts). Future research, building upon these theoretical frameworks, is needed to determine the ways in which gender (and other moderators) interact with sexual abuse in relation to particular mediating processes and specific psychological outcomes.

There are several limitations represented in the current state of the field that may have influenced our ability to detect patterns of specificity between particular stressors and broad-band outcomes. First, there are significant rates of co-occurrence and comorbidity of psychopathology among youth (Compas & Hammen, 1994). Second, there are difficulties associated with the lack of a well-established

taxonomy of stressors. Categorizing stressors based on type may include a variety of stressors that were different in nature. For example, poverty is associated with a range of additional specific stressful experiences, which include both 'conflict' (e.g., exposure to violence, physical abuse, divorce/marital conflict) and 'loss'-based stressors (e.g., deaths, separations, disasters). A third explanation is that specificity findings are moderated by specific child-based characteristics and environment-based contexts. Thus, particular stressors may be linked with particular outcomes, only in the presence of particular moderating and mediating processes.

It should also be noted that given the variability in methodology, stressors, outcomes, measures, and samples, we determined that a qualitative approach to the review would be the most appropriate; however, as the field of developmental psychopathology progresses, a meta-analytic approach to reviewing the literature is recommended. Further, although variability in measurement and definition led us to focus on internalizing versus externalizing outcomes, it is possible that this more global review led to more difficulty in detecting evidence of specificity, which, by definition, necessitates examining specific stressors in relation to specific outcomes. In addition to these limitations, there are several methodological issues that should be considered within the context of this review.

### *Methodological issues*

Specificity research has been fraught with variability in specific constructs examined, measures used, and samples studied, seriously limiting conclusions which may be drawn about specificity itself and conclusions about why discrepancies in findings exist. Further, specific methods for examining specificity have varied. Beyond the level of design (Stressor Specific, Outcome Specific, and Stressor-Outcome Specific), analytical and statistical procedures utilized were found to be quite disparate across studies. For instance, some studies utilized a 'comparison approach' in which rates of psychological problems were compared across groups exposed to differing stressors; whereas other studies utilized a 'relational approach' in which associations between particular stressors and particular outcomes were examined in a single sample of youth. Thus, the approach utilized is likely to influence the findings reported. Although there is considerable heterogeneity within these approaches, the relational approach is generally more stringent than the comparison approach.

Most of the research conducted in the stress and coping literature is cross-sectional in nature. Thus, it is unclear the extent to which relations between stressors and outcomes change across time. A developmental approach, which takes into account changes across time and developmental trends, to

understanding specificity, equifinality, and multifinality may improve our understanding of the pathways between stress and psychopathology. For example, the way children express their distress in response to stressors changes over the course of development, so age may be one potential explanation for the mixed findings on specificity.

It may also be helpful to consider differences in sampling strategies (i.e., clinical samples versus population-based samples) in examining specificity. For example, comorbidity of mood and conduct disorders in children and adolescents has been found to be higher in clinic samples than community populations. Thus, it is possible that greater specificity may exist between particular stressors and particular outcomes in community-based samples than clinic samples. In addition, different theoretical frameworks and methodological characteristics of studies may pull for particular specificity results. For example, links between parent reports of abuse and child externalizing symptoms may be stronger due to mono-informant or mono-method bias or the difficulty of assessing internalizing symptoms in a psychometrically sound manner.

Related to methodological variability is the dearth of theory-driven specificity research. In light of the large number of studies meeting minimal methodological requirements for specificity research, it is striking how few studies tested theory-based models of specificity. The work of Sandler and colleagues (1992) and Shirk and colleagues (1998) represent important exceptions. More studies like these, which test specific etiological models of developmental psychopathology, are needed.

### **Directions for future research**

There is a need to develop and test specific models that examine the complex relations between stress and psychopathology. It is noteworthy that no single study has yet tested full specificity models (including examination of specific mediating and moderating pathways) of the role of stress in the etiology of a particular psychological outcome (see Figure 2). For example, exposure to conflict among adolescent boys leads to external, global, stable cognitive appraisals, which lead to aggression symptoms, which, in turn, lead to further conflict-based stressors. Such research is necessary to develop an empirically supported model of the role of stress in the etiology of developmental psychopathology.

Further discussion of the theoretical model presented (see Figure 2) may help to illustrate and highlight the type of work that is necessary in order to better understand the complex relations between stress and psychological outcomes. This model may be tested with various simple and complex, theoretically driven hypotheses regarding relationships between stressors and outcomes. For an example,

research testing such a model might 1) examine the influence of a particular stressor to a particular outcome (e.g., test the hypothesis that there is an association between exposure to violence and aggressive behavior by using the 'relational' approach of examining several stressors – i.e., exposure to violence, poverty, and divorce, and several outcomes – i.e., aggressive behavior, depressive symptoms and anxiety symptoms), or 2) examine both the specificity of the association between exposure to violence and aggressive behavior (as in example 1) and the influence of a particular moderator on the relation between a particular stressor and a particular outcome (e.g., test the hypothesis that male gender strengthens the association between exposure to violence and aggressive behavior), or 3) examine specificity (example 1), a potential moderator (example 2), and the influence of a particular moderator on both a particular stressor and a particular mediator (e.g., test the hypothesis that male gender strengthens the association between exposure to violence and aggressive behavior through increases in distraction and avoidant coping). In addition, reciprocal and dynamic relations among a particular moderator and a particular stressor, outcome, and/or mediating process could be examined. For example, the hypothesis that psychological problems (e.g., aggressive behavior) lead to the development of a moderating context (e.g., hostility from classmates at school) that, in turn, exacerbates the association between a particular stressor (e.g., a violent attack at school) and additional specific psychological symptoms might be tested.

In order to begin to establish more comparability between studies and to conduct a clear test of specificity, it is recommended that researchers consider the following criteria: 1) test specific, theory-based, etiological models of developmental psychopathology; 2) include at least two stressors and at least two outcomes in order to examine specificity with the more stringent Stressor-Outcome model; 3) develop a taxonomy of stressors, and in the meantime, consider refining notions of categorizing stressful life events based on theoretically-derived categories (i.e., conflict, loss, achievement); 4) identify both unique and common effects of various stressors in relation to various outcomes; 5) include moderators, such as child characteristics and environmental context, as well as mediators, such as cognitions and coping, in the examination of specificity; 6) use a relational approach, examining the relations between stressors and outcomes within a particular population; 7) use multiple methods of assessment to reduce reporter bias and generate more confidence in the findings; and 8) conduct longitudinal studies to examine the dynamic relations between stressors and outcomes across time.

In sum, methodological discrepancies and the lack of theory-driven research have limited the degree to which the specificity hypothesis has been tested.

Although the results of the present review provide little evidence of specificity effects, examination of full specificity models (using maximally effective methodology) is needed to fully test hypotheses related to equifinality, multifinality, and specificity in the relation between stress and psychological symptoms in children and adolescents. Results of such research may, in fact, provide evidence that integrates these theories (i.e., through discovery that specific relations between particular stressors and particular outcomes occur only in the context of specific moderators and/or specific mediating processes). Such research provides an ideal method for examining the individual patterns of adaptive and maladaptive outcomes across development (Cicchetti & Rogosch, 1996) and, thus, furthering the goals of developmental psychopathology.

### Correspondence to

Susan D. McMahon, Department of Psychology, DePaul University, 2219 N. Kenmore, Chicago, IL, 60614, USA; Tel: 773-325-2039; Email: smcmahon@depaul.edu

### References

- Achenbach, T.M. (1991). Comorbidity in child and adolescent psychiatry: Categorical and quantitative perspectives. *Journal of Child and Adolescent Psychopharmacology*, 1, 271–278.
- Achenbach, T.M., & Howell, C.T. (1993). Are American children's problems getting worse? *Journal of the American Academy of Child and Adolescent Psychiatry*, 32, 1145–1154.
- Achenbach, T.M., & Dumenci, L. (2001). Advances in empirically based assessment: Revised cross-informant syndromes and new DSM-oriented scales for the CBCL, YSR, and TRF: Comment on Lengua, Sadowksi, Friedrich, and Fisher (2001). *Journal of Consulting and Clinical Psychology*, 69, 699–702.
- Ackerman, P.T., Newton, J.E.O., McPherson, W.B., Jones, J.G., & Dykman, R.A. (1998). Prevalence of post traumatic stress disorder and other psychiatric diagnoses in three groups of abused children (sexual, physical, and both). *Child Abuse and Neglect*, 22, 759–774.
- Aseltine, R., & Gore, S. (1993). Mental health and social adaptation following the transition from high school. *Journal of Research on Adolescence*, 3, 347–370.
- Attar, B., Guerra, N., & Tolan, P. (1994). Neighborhood disadvantage, stressful life events, and adjustment in urban elementary-school children. *Journal of Clinical Child Psychology*, 23, 391–400.
- Bayatpour, M., Wells, R., & Holford, S. (1992). Physical and sexual abuse as predictors of substance use and suicide among pregnant teenagers. *Journal of Adolescent Health*, 13, 128–132.
- Berliner, L., & Elliott, D.M. (1996). Sexual abuse of children. In J. Briere, L. Berliner, J.A. Bulkley, C. Jenny, & T. Reid (Eds.), *The APSAC handbook on*

- child maltreatment* (pp. 51–71). Thousand Oaks: Sage Publications.
- Berliner, L., & Conte, J. (1995). The effects of disclosure and intervention on sexually abused children. *Child Abuse and Neglect, 19*, 371–384.
- Black, M., Dubowitz, H., & Harrington, D. (1994). Sexual abuse: Developmental differences in children's behavior and self-perception. *Child Abuse and Neglect, 18*, 85–95.
- Bolger, K.E., Patterson, C.J., Thompson, W.W., & Kupersmidt, J.B. (1995). Psychosocial adjustment among children experiencing persistent and intermittent family economic hardship. *Child Development, 66*, 1107–1129.
- Boney-McCoy, S., & Finkelhor, D. (1995). Psychosocial sequelae of violent victimization in a national youth sample. *Journal of Consulting and Clinical Psychology, 63*, 726–736.
- Brand, E.F., King, C.A., Olson, E., Ghazziuddin, N., & Naylor, M. (1996). Depressed adolescents with a history of sexual abuse: Diagnostic comorbidity and suicidality. *Journal of the American Academy of Child and Adolescent Psychiatry, 35*, 34–41.
- Brent, D., Moritz, G., Bridge, J., Perper, J., & Canobbio, R. (1996). The impact of adolescent suicide on siblings and parents: A longitudinal follow-up. *Suicide and Life-Threatening Behavior, 26*, 253–259.
- Brent, D., Perper, J., Moritz, G., Friend, A., Schweers, J., Allman, C., McQuiston, L., Boylan, M.B., Roth, C., & Balach, L. (1993). Adolescent witness to a peer suicide. *Journal of the American Academy of Child and Adolescent Psychiatry, 32*, 1184–1188.
- Brent, D., Perper, J., Moritz, G., Allman, C., Schweers, J., Roth, C., Balach, L., Canobbio, R., & Liotus, L. (1993). Psychiatric sequelae to the loss of an adolescent peer to suicide. *Journal of the American Academy of Child and Adolescent Psychiatry, 32*, 509–517.
- Breton, J.J., Valla, J.P., & Lambert, J. (1993). Industrial disaster and mental health of children and their parents. *Journal of Academic Child Adolescent Psychiatry, 32*, 438–445.
- Brody, G., & Forehand, R. (1990). Interparental conflict, relationship with the noncustodial father, and adolescent post-divorce adjustment. *Journal of Applied Developmental Psychology, 11*, 139–147.
- Brody, G., & Flor, D. (1997). Maternal psychological functioning, family processes, and child adjustment in rural, single-parent, African American families. *American Psychological Association, 6*, 1000–1011.
- Burke, P., Meyer, V., Kocoshis, S., Orenstein, D., Chandra, R., Nord, D., Sauer, J., & Cohen, E. (1989). Depression and anxiety in pediatric inflammatory bowel disease and cystic fibrosis. *Journal of the American Academy of Child and Adolescent Psychiatry, 28*, 948–951.
- Caviola, A., & Schiff, M. (1988). Behavioral sequelae of physical and/or sexual abuse in adolescents. *Child Abuse and Neglect, 12*, 181–188.
- Chandy, J.M., Blum, R.W., & Resnick, M.D. (1996). Gender-specific outcomes for sexually abused adolescents. *Child Abuse and Neglect, 20*, 1219–1231.
- Children's Defense Fund (1999). *The state of America's children yearbook*. Washington, DC: Children's Defense Fund.
- Cicchetti, D., & Cohen, D.J. (1995). Perspectives on developmental psychopathology. In D. Cicchetti & D.J. Cohen (Eds.), *Developmental psychopathology volume 1: Theory and methods* (pp. 3–20). New York: John Wiley & Sons.
- Cicchetti, D., & Toth, S.L. (1991). A developmental perspective on internalizing and externalizing disorders. In D. Cicchetti & S.L. Toth (Eds.), *Internalizing and externalizing expressions of dysfunction* (pp. 1–19). New York: Erlbaum.
- Cicchetti, D., & Toth, S.L. (Eds.). (1997). *Developmental perspectives on trauma: Theory, research and intervention*. Rochester, NY: Rochester University Press.
- Cicchetti, D., & Rogosch, F.A. (1996). Equifinality and multifinality in developmental psychopathology. *Development and Psychopathology, 8*, 597–600.
- Cohen, J.A., & Mannarino, A.P. (1988). Psychological symptoms in sexually abused girls. *Child Abuse and Neglect, 12*, 571–577.
- Cohen, L.H., & Park, C. (1992). Life stress in children and adolescents: An overview of conceptual and methodological issues. In A.M. LaGraca, L.J. Siegel, J.L. Wallander, & C.E. Walker (Eds.), *Stress and coping in child health* (pp. 25–43). New York: Guilford.
- Cohen, Y., Spirito A., Sterling C., Donaldson D., Seifer, R., Plummer, B., Avila, R., & Ferrer, K. (1996). Physical and sexual abuse and their relation to psychiatric disorder and suicidal behavior among adolescents who are psychiatrically hospitalized. *Journal of Child Psychology and Psychiatry, 37*, 989–993.
- Compas, B.E. (1987). Stress and life events during childhood and adolescence. *Clinical Psychology Review, 7*, 275–302.
- Compas, B.E., & Hammen, C.L. (1994). Child and adolescent depression: Covariation and comorbidity in development. In R.J. Haggerty, N. Garnezy, M. Rutter, & L. Sherrrod, (Eds.), *Risk and resilience in children: Developmental approaches* (pp. 225–267). New York: Cambridge University Press.
- Conger, R.D., Ge, X., Elder, G.H., Jr., Lorenz, F.O., & Simons, R.L. (1994). Economic stress, coercive family process, and developmental problems of adolescents. *Child Development, 65*, 541–561.
- Conger, R.D., Conger, K.J., Elder, G.H., Jr., Lorenz, F., Simons, R., & Whitbeck, L. (1993). A family process model of economic hardship and adjustment of early adolescent girls. *Developmental Psychology, 29*, 206–219.
- Conger, R.D., Conger, K.J., Elder, G.H., Jr., Lorenz, F., Simons, R., & Whitbeck, L. (1992). A family process model of economic hardship and adjustment of early adolescent boys. *Child Development, 63*, 526–541.
- Cooley-Quille, M.R., Turner, S.M., & Beidel, D.C. (1995). Emotional impact of children's exposure to community violence: A preliminary study. *Journal of the American Academy of Child and Adolescent Psychiatry, 34*, 1362–1368.
- Cosentino, C.E., Meyer-Bahlburg, H.F.L., Alpert, J.L., Weinberg, S.L., & Gaines, R. (1995). Sexual behavior problems and psychopathology symptoms in sexually abused girls. *Journal of the American Academy of Child and Adolescent Psychiatry, 34*, 1033–1042.
- Crittenden, P., Claussen, A., & Sugarman, D. (1994). Physical and psychological maltreatment in middle

- childhood and adolescence. *Development and Psychopathology*, 6, 145–164.
- David, C., Steel, R., Forehand, R., & Armistead, L. (1996). The role of family conflict and marital conflict in adolescent functioning. *Journal of Family Violence*, 1, 81–91.
- Deblinger, E., McLeer, S., Atkins, M., Ralphe, D., & Foa, E. (1989). Post-traumatic stress in sexually abused, physically abused, and nonabused children. *Child Abuse and Neglect*, 13, 403–408.
- DePaul, J., & Arruabarrena, M.I. (1995). Behavior problems in school-aged physically abused and neglected children in Spain. *Child Abuse and Neglect*, 19, 409–418.
- DuBois, D., Felner, R., Meares, H., & Krier, M. (1994). Prospective investigation of the effects of socioeconomic disadvantage, life stress, and social support on early adolescent adjustment. *Journal of Abnormal Psychology*, 103, 511–522.
- Durant, R.H., Pendergrast, R.A., & Cadenhead, C. (1994). Exposure to violence and victimization and fighting behavior by urban black adolescents. *Journal of Adolescent Health*, 15, 311–318.
- Durkin, M.S., Khan, N., Davidson, L.L., Zaman, S.S., & Stein, Z.A. (1993). The effects of a natural disorder on child behavior: Evidence for posttraumatic stress. *American Journal of Public Health*, 83, 1549–1553.
- Dykman, R.A., McPherson, B., Ackerman, P.T., Newton, J.E.O., Mooney, D.M., Wherry, J., & Chaffin, M. (1997). Internalizing and externalizing characteristics of sexually and/or physically abused children. *Integrative Physiological and Behavioral Science*, 32, 62–83.
- Einbender, A., & Friedrich, W.N. (1989). Psychological functioning and behavior of sexually abused girls. *Journal of Consulting and Clinical Psychology*, 57, 155–157.
- Elder, G.H., Conger, R.D., Foster, M.E., & Ardel, M. (1992). Families under economic pressure. *Journal of Family Issues*, 13, 5–37.
- Engstrom, I. (1992). Mental health and psychological functioning in children and adolescents with Inflammatory Bowel Disease: A comparison with children with children having other chronic illnesses and with healthy children. *Journal of Child Psychology and Psychiatry*, 33, 563–602.
- Famularo, R., Kinscherff, R., & Fenton, T. (1992). Psychiatric diagnoses of maltreated children: Preliminary findings. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31, 863–867.
- Feldman, R.S., Salzinger, S., Rosario, M., Alvarado, L., Caraballo, L., & Hammer, M. (1995). Parent, teacher, and peer ratings of physically abused and maltreated children's behavior. *Journal of Abnormal Child Psychology*, 23, 317–335.
- Felner, R.D., Brand, S., DuBois, D.L., Adan, A., Mulhall, P., & Evans, E. (1995). Socioeconomic disadvantage, proximal environmental experiences, and socioemotional and academic adjustment in early adolescence: Investigation of a mediated effects model. *Child Development*, 66, 774–792.
- Fergusson, D.M., Horwood, L.J., & Lynskey, M. (1994). Parental separation, adolescent psychopathology, and problem behaviors. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33, 1122–1131.
- Finkelhor, D., & Brown, A. (1985). The traumatic impact of child sexual abuse: A conceptualization. *American Journal of Orthopsychiatry*, 55, 530–541.
- Flisher, A.J., Kramer, R.A., Hoven, C.W., & Greenwald, S. (1997). Psychosocial characteristics of physically abused children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36, 123–131.
- Forehand, R., McCombs, A., Long, N., Brody, G., & Fauber, R. (1988). Early adolescent adjustment to recent parental divorce: The role of interparental conflict and adolescent sex as mediating variables. *Journal of Consulting and Clinical Psychology*, 56, 624–627.
- Forehand, R., Neighbors, B., & Wierson, M. (1991). The transition to adolescence: The role of gender and stress in problem behavior and competence. *Association for Child Psychology and Psychiatry*, 6, 929–937.
- Freeman, L., Shaffer, D., & Smith, H. (1996). Neglected victims of homicide: The needs of young siblings of murder victims. *American Journal of Orthopsychiatry*, 66, 337–345.
- Friedrich, W.N., Jaworski, T.M., Huxsahl, J.E., & Bengtson, B.S. (1997). Dissociative and sexual behaviors in children and adolescents with sexual abuse and psychiatric histories. *Journal of Interpersonal Violence*, 12, 155–171.
- Gale, J., Thompson, R.J., Moran, T., & Sack, W.H. (1988). Sexual abuse in young children: Its clinical presentation and characteristics patterns. *Child Abuse and Neglect*, 12, 163–170.
- Garber, J., & Hollon, S. (1991). What can specificity designs say about causality in psychopathology research? *Psychological Bulletin*, 110, 129–136.
- Garnezy, N. (1987). Stress, competency and development: Continuities in the study of schizophrenic adults, children vulnerable to psychopathology, and the search for stress-resistant children. *American Journal of Orthopsychiatry*, 57, 159–174.
- Garnefski, N.Y., & Arends, E. (1998). Sexual abuse and adolescent maladjustment: Differences between male and female victims. *Journal of Adolescence*, 21, 99–107.
- Garralda, M.E., Jameson, R.A., Reynolds, J.M., & Postlethwaite, R.J. (1988). Psychiatric adjustment in children with chronic renal failure. *Journal of Child Psychology and Psychiatry*, 29, 79–90.
- Glod, C.A., & Teicher, M.H. (1996). Relationship between early abuse, posttraumatic stress disorder, and activity levels in prepubertal children. *Journal of the American Academy of Child and Adolescent Psychiatry*, 35, 1384–1393.
- Goldston, D.B., Turnquist, D.C., & Knutson, J.F. (1989). Presenting problems of sexually abused girls receiving psychiatric services. *Journal of Abnormal Psychology*, 98, 314–317.
- Gorman-Smith, D., & Tolan, P. (1998). The role of exposure to community violence and developmental problems among inner-city youth. *Development and Psychopathology*, 10, 101–116.
- Gould, M.S., Shaffer, D., Fisher, P., & Garfinkel, R. (1998). Separation/divorce and child and adolescent

- completed suicide. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37, 155–162.
- Grant, K.E., Compas, B.E., Stuhlmacher, A.F., Thurm, A.E., McMahon, S.D., & Halpert, J.A. (in press). Stressors and child/adolescent psychopathology: Moving from markers to mechanisms of risk. *Psychological Bulletin*.
- Haggerty, R.J., Sherrod, L.R., Garmezy, N., & Rutter, M. (Eds.). (1994). *Stress, risk and resilience in children and adolescents: Processes, mechanisms, and interventions*. New York: Cambridge University Press.
- Haviland, M.G., Sonne, J.L., & Woods, L.R. (1995). Beyond posttraumatic stress disorder: Object relations and reality testing disturbances in physically and sexually abused adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34, 1054–1059.
- Hazzard, A., Celano, M., Gould, J., Lawry, S., & Webb, C. (1995). Predicting symptomatology and self-blame among child sex abuse victims. *Child Abuse and Neglect*, 19, 707–714.
- Hennessy, K.D., Rabideau, G.J., Cicchetti, D., & Cummings, E.M. (1994). Responses of physically abused and nonabused children to different forms of interadult anger. *Child Development*, 65, 815–828.
- Hernandez, J., Lodico, M., & DiClemente, R. (1993). The effects of child abuse and race on risk-taking in male adolescents. *Journal of the National Medical Association*, 85, 593–597.
- Hernandez, J. (1995). The concurrence of eating disorders with histories of child abuse among adolescents. *Journal of Child Sexual Abuse*, 4, 73–85.
- Hibbard, R.A., & Hartman, G.L. (1992). Behavioral problems in alleged sexual abuse victims. *Child Abuse and Neglect*, 16, 755–762.
- Higgins, D.J., & McCabe, M.P. (1998). Parent perceptions of maltreatment and adjustment in children. *Journal of Family Studies*, 4, 53–76.
- Hoyt, L.A., Cowen, E.L., Pedro-Carroll, J.L., & Alpert-Gillis, L.J. (1990). Anxiety and depression in young children of divorce. *Journal of Clinical Child Psychology*, 19, 26–32.
- Hussey, D.L., & Singer, M. (1993). Psychological distress, problem behaviors, and family functioning of sexually abused adolescent inpatients. *Journal of Academic Child Adolescent Psychiatry*, 32, 954–961.
- Hussey, D.L., Strom, G., & Singer, M. (1992). Male victims of sexual abuse: An analysis of adolescent psychiatric inpatients. *Child and Adolescent Social Work Journal*, 9, 491–503.
- Inderbitzen-Pisaruk, H., Shawchuck, C.R., & Hoier, T. (1992). Behavioral characteristics of child victims of sexual abuse: A comparison study. *Journal of Clinical Child Psychology*, 21, 14–19.
- Jarvelin, M.R., Moilanen, I., Vikevainen-Tervonen, L., & Huttunen, N.P. (1990). Life changes and protective capacities in enuretic and non-enuretic children. *Journal of Child Psychology and Psychiatry*, 31, 763–74.
- Jenkins, J.M., & Smith, M.A. (1990). Factors protecting children living in disharmonious homes: Maternal reports. *Journal of the American Academy of Child and Adolescent Psychiatry*, 29, 60–69.
- Jenkins, J.M., & Smith, M.A. (1991). Marital disharmony and children's behavior problems: Aspects of a poor marriage that affect children adversely. *Journal of Child Psychology and Psychiatry*, 32, 793–810.
- Johnson, J.H. (1986). *Life events as stressors in childhood and adolescence*. Beverly Hills: Sage Publications.
- Johnson, J.H., & Bradlyn, A.S. (1988). Assessing stressful life events in childhood and adolescence. In P. Karoly (Ed.), *Handbook of child health assessment: Biopsychosocial perspectives* (pp. 303–331). New York: John Wiley & Sons.
- Johnson, B.K., & Kenkel, M.B. (1991). Stress, coping, and adjustment in female adolescent incest victims. *Child Abuse and Neglect*, 15, 293–305.
- Johnston, J.R., Gonzalez, R., & Campbell, L.E. (1987). Ongoing postdivorce conflict and child disturbance. *Journal of Abnormal Child Psychology*, 15, 493–509.
- Jouriles, E.N., Norwood, W.D., McDonald, R., Vincent, J.P., & Mahoney, A. (1996). Physical violence and other forms of marital aggression: Links with children's behavior problems. *Journal of Family Psychology*, 10, 223–234.
- Jouriles, E.N., Barling, J., & O'Leary, K.D. (1987). Predicting child behavior problems in maritally violent families. *Journal of Abnormal Child Psychology*, 15, 165–173.
- Kaplan, S.J., Pelcovitz, D., Salzinger, S., Mandel, F., & Weiner, M. (1997). Adolescent physical abuse and suicide attempts. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36, 799–808.
- Kashani, J.H., Konig, P., Shepperd, J.A., Wilfley, D., & Morris, D.A. (1988). Psychopathology and self-concept in asthmatic children. *Journal of Pediatric Psychology*, 13, 509–520.
- Kolko, D.J., Moser, J., & Weldy, S.R. (1988). Behavioral/emotional indicators of sexual abuse in child psychiatric inpatient: A controlled comparison with physical abuse. *Child Abuse and Neglect*, 12, 529–541.
- Kolko, D.J., & Kazdin, A.E. (1993). Emotional/Behavioral problems in clinic and nonclinic children: Correspondence among child, parent and teacher reports. *Journal of Child Psychology and Psychiatry*, 34, 991–1006.
- Kuhn, J.A., Arellano, C.M., & Chavez, E.L. (1998). Correlates of sexual assault in Mexican American and white non-Hispanic adolescent males. *Violence and Victims*, 13, 11–20.
- Kumar, G., Steer, R.A., & Deblinger, E. (1996). Problems in differentiating sexually from nonsexually abused adolescent psychiatric inpatients by self-reported anxiety, depression, internalization, and externalization. *Journal of Child Abuse and Neglect*, 20, 1070–1086.
- La Greca, A.M., Silverman, W.K., & Wasserstein, S.B. (1998). Children's predisaster functioning hurricane Andrew. *Journal of Consulting and Clinical Psychology*, 66, 883–892.
- Laor, N., Wolmer, L., Mayes, L.C., Golomb, A., Silverberg, D.S., Weizman, R., & Cohen, D. (1996). Israeli preschoolers under scud missile attacks: A developmental perspective on risk-modifying factors. *Archives of General Psychiatry*, 53, 416–423.



- Lazarus, R.S., & Folkman, S. (1984). *Stress, appraisal and coping*. New York: Springer.
- Lempers, J., Clark-Lempers, D., & Simons, R. (1989). Economic hardship, parenting practices, and adolescent distress. *Child Development, 60*, 25–39.
- Levendosky, A.A., & Graham-Bermann, S.A. (1998). The moderating effects of parenting stress on children's adjustment in woman-abusing families. *Journal of Interpersonal Violence, 13*, 383–397.
- Lewinsohn, P.M., Gotlib, I.H., & Seeley, J.R. (1995). Adolescent psychopathology: IV. Specificity of psychosocial risk factors for depression and substance abuse in older adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry, 34*, 1221–1229.
- Livingston, R., Lawson, R., & Jones, J. (1993). Predictors of self-reported psychopathology in children abused repeatedly by a parent. *Journal of the American Academy of Child and Adolescent Psychiatry, 32*, 948–953.
- Long, N., Slater, E., Forehand, R., & Fauber, R. (1988). Continued high or reduced interparental conflict following divorce: Relation to young adolescent adjustment. *Journal of Consulting and Clinical Psychology, 56*, 467–469.
- Luster, T., & Small, S.A., (1997). Sexual abuse history and problems in adolescence: Exploring the effects of moderating variables. *Journal of Marriage and the Family, 59*, 131–142.
- Lynch, M., & Cicchetti, D. (1998). An ecological-transactional analysis of children and contexts: The longitudinal interplay among child maltreatment, community violence, and children's symptomatology. *Development and Psychopathology, 10*, 235–257.
- Manly, J.T., Cicchetti, D., & Barnett, D. (1994). The impact of subtype, frequency, chronicity and severity of child maltreatment on social competence and behavior problems. *Development and Psychopathology, 6*, 121–143.
- Mannarino, A.P. (1988). Psychological symptoms exhibited by children who had recently been sexually abused. *Journal of Child Abuse and Neglect, 12*, 571–577.
- Mannarino, A.P., & Cohen, J.A. (1996). Abuse-related attributions and perceptions, general attributions, and locus of control in sexually abused girls. *Journal of Interpersonal Violence, 11*, 162–180.
- March, J.S., Amaya-Jackson, L., Terry, R., & Costanzo, P. (1997). Posttraumatic symptomatology in children and adolescents after an industrial fire. *Journal of the American Academy of Child and Adolescent Psychiatry, 36*, 1080–1088.
- Markward, M.J. (1997). The impact of domestic violence on children. *Society, 78*, 66–70.
- Martin, G. (1996). Reported family dynamics, sexual abuse, and suicidal behaviors in community adolescents. *Archives of Suicide Research, 2*, 183–195.
- Marttunen, M.J., Aro, H.M., Henriksson, M.M., & Loennqvist, J.K. (1994). Psychosocial stressors more common in adolescent suicides with alcohol abuse compared with depressive adolescent suicides. *Journal of the American Academy of Child and Adolescent Psychiatry, 33*, 490–497.
- Masten, A.S., Miliotis, D., Graham-Berman, S.A., Ramirez, M., & Neemann, J. (1993). Children in homeless families: Risks to mental health and development. *Journal of Consulting and Clinical Psychology, 61*, 335–343.
- McClellan, J., Adams, J., Douglas, D., McCurry, C., & Storck, M. (1995). Clinical characteristics related to severity of sexual abuse: A study of seriously mentally ill youth. *Child Abuse and Neglect, 19*, 1245–1254.
- McCloskey, L.A., Figueredo, A. Jose, & Koss, M.P. (1995). The effects of systemic family violence on children's mental health. *Child Development, 66*, 1239–1261.
- McLeer, S., Callaghan, J., Henry, D., & Wallen, J. (1994). Psychiatric disorders in sexually abused children. *Journal of the American Academy of Child and Adolescent Psychiatry, 33*, 313–319.
- McLeer, S., Deblinger, E., Henry, D., & Orvaschel, H. (1992). Sexually abused children at high risk for post-traumatic stress disorder. *Journal of the American Academy Child and Adolescent Psychiatry, 31*, 875–879.
- McLoyd, V.C., Jayaratne, T.E., Beballo, R., & Borquez, J. (1994). Unemployment and work interruption among African American single mothers: Effects on parenting and adolescent socioemotional functioning. *Child Development, 65*, 562–589.
- Mezzich, A.C., Giancola, P.R., Tarter, R.E., Lu, S., Parks, S.M., & Barrett, C.M. (1997). Violence, suicidality, and alcohol/drug use involvement in adolescent females with a psychoactive substance use disorder and controls. *Alcoholism, Clinical and Experimental Research, 21*, 1300–1307.
- Mian, M., Marton, P., & LeBaron, D. (1996). The effects of sexual abuse on 3- to 5- year-old girls. *Journal of Child Abuse and Neglect, 20*, 731–745.
- Mikail, S.F., & Von Baeyer, C.L. (1990). Pain, somatic focus, and emotional adjustment in children of chronic headache sufferers and controls. *Social Science Medicine, 31*, 51–59.
- Murata, J. (1994). Family stress, social support, violence, and sons' behavior. *Western Journal of Nursing Research, 16*, 154–168.
- Najarian, L., Goenjian, A., Pelcovitz, D., Mandel, F., & Najarian, B. (1996). Relocation after a disaster: Posttraumatic stress disorder in Armenia after the earthquake. *Journal of the American Academy of Child and Adolescent Psychiatry, 35*, 374–383.
- O'Keefe, M. (1994a). Linking marital violence, mother-child/father-child aggression, and child behavior problems. *Journal of Family Violence, 9*, 63–78.
- O'Keefe, M. (1994b). Racial/ethnic differences among battered women and their children. *Journal of Child and Family Studies, 3*, 283–305.
- O'Keefe, M. (1997). Adolescents' exposure to community and school violence: Prevalence and behavioral correlates. *Journal of Adolescent Health, 20*, 368–376.
- O'Koon, J. (1999). *Covariation of internalizing and externalizing symptoms among low-income urban youth*. Unpublished doctoral dissertation, DePaul University, Chicago.
- Osofsky, J.D., Wewers, S., Hann, D.M., & Fick, A.C. (1993). Chronic community violence: What is happening to our children? *Psychiatry, 56*, 36–45.

- Pelcovitz, D., Kaplan, S., Goldenberg, B., Mandel, F., Lehane, J., & Guarrera, J. (1994). Post-traumatic stress disorder in physically abused adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry, 33*, 305-312.
- Pfeffer, C.R., Martins, P., Mann, J., Sunkenberg, M., Ice, A., Damore, J., Gallo, C., Karpenos, I., & Jiang, H. (1997). Child survivors of suicide: Psychosocial characteristics. *Journal of the American Academy of Child and Adolescent Psychiatry, 36*, 65-74.
- Prino, C., & Peyrot, M. (1994). The effect of child abuse and neglect on aggressive, withdrawn, and prosocial behavior. *Child Abuse and Neglect, 18*, 871-884.
- Reese, F.L., & Roosa, M.W. (1991). Early adolescents' self-reports of major life stressors and mental health risk status. *Journal of Early Adolescence, 11*, 363-378.
- Reinherz, H., Giaconia, R., Pakiz, B., Silverman, A., Frost, A., & Lefkowitz, E. (1993). Psychosocial risks for major depression in late adolescence: A longitudinal community study. *Journal of the American Academy of Child and Adolescent Psychiatry, 32*, 1155-1163.
- Rivinus, T., Levoy, D., Matzko, M., & Seifer, R. (1992). Hospitalized children of substance-abusing parents and physically and sexually abused children: A comparison. *Journal of the American Academy of Child and Adolescent Psychiatry, 31*, 1019-1023.
- Rodgers, B. (1994). Pathways between parental divorce and adult depression. *Journal of Child Psychology and Psychiatry, 35*, 1289-1308.
- Rossmann, B.B.R., Bingham, R.D., & Emde, R.N. (1997). Symptomatology and adaptive functioning for children exposed to normative stressors, dog attack, and parental violence. *Journal of the American Academy of Child and Adolescent Psychiatry, 36*, 1089-1097.
- Rotheram-Borus, M.J., Mahler, K.A., Koopman, C., & Langabeer, K. (1996). Sexual abuse history and associated multiple risk behavior in adolescent run-aways. *American Journal of Orthopsychiatry, 66*, 390-400.
- Rutter, M. (1989). Pathways from childhood to adult life. *Journal of Child Psychology and Psychiatry, 30*, 23-51.
- Sadeh, A., Hayden, R.M., McGuire, J.P.D., Sachs, H., & Civita, R. (1994). Somatic, cognitive and emotional characteristics of abused children in a psychiatric hospital. *Child Psychiatry and Human Development, 24*, 191-199.
- Sandler, I.N., Reynolds, K.D., Kliever, W., & Ramirez, R. (1992). Specificity of the relation between life events and psychological symptomatology. *Journal of Clinical Child Psychology, 21*, 240-248.
- Schwab-Stone, M., Ayers, T., Kaspro, W., Voyce, C., Barone, C., Shriver, T., & Weissberg, R. (1995). No safe haven: A study of violence exposure in an urban community. *Journal of the American Academy of Child and Adolescent Psychiatry, 34*, 1343-1352.
- Selye, H. (1956). *The stress of life*. New York: McGraw-Hill.
- Selye, H. (1974). *Stress without distress*. Philadelphia: Lippincott.
- Shapiro, J.P., Leifer, M., Martone, M.W., & Kassem, L. (1992). Cognitive functioning and social competence as predictors of maladjustment in sexually abused girls. *Journal of Interpersonal Violence, 7*, 156-164.
- Shaw, J.A., Applegate, B., Tanner, S., Perez, D., Rothe, E., Campo-Bowen, A.E., & Lahey, B.L. (1995). Psychological effects of Hurricane Andrew on an elementary school population. *American Academy of Child and Adolescent Psychiatry, 34*, 1185-1192.
- Shirk, S., Boergers, J., Eason, A., & Van Horn, M. (1998). Dysphoric interpersonal schemata and pre-adolescents' sensitization to negative events. *Journal of Clinical Child Psychology, 27*, 54-65.
- Sibthorpe, B., Drinkwater, J., Gardner, K., & Bammer, G. (1995). Drug use, binge drinking and attempted suicide among homeless and potentially homeless youth. *Australian and New Zealand Journal of Psychiatry, 29*, 248-256.
- Siegel, K., Mesagno, F., Karus, D., Christ, G., Banks, K., & Moynihan, R. (1992). Psychosocial adjustment of children with a terminally ill parent. *Journal of the American Academy of Child and Adolescent Psychiatry, 31*, 327-333.
- Silverman, A.B., Reinherz, H.Z., & Giaconia, R.M. (1996). The long-term sequelae of child and adolescent abuse: A longitudinal community study. *Child Abuse and Neglect, 20*, 709-723.
- Smith, S.L., & Howard, J.A. (1994). The impact of previous sexual abuse on children's adjustment in adoptive placement. *Social Work, 39*, 491-501.
- Smith, S.L., Howard, J.A., & Monroe, A.D. (1998). An analysis of child behavior problems in adoptions in difficulty. *Journal of Social Research, 24*, 61-84.
- Spaccarelli, S., Sandler, I.N., & Roosa, M. (1994). History of spouse violence against mother: Correlated risks and unique effects in child mental health. *Journal of Family Violence, 9*, 79-98.
- Spaccarelli, S. (1994). Stress appraisal and coping in child sexual abuse: A theoretical and empirical review. *Psychological Bulletin, 116*, 340-362.
- Stern, A.E., Lynch, D.L., Oates R.K., O'Toole, B.I., & Cooney, G. (1995). Self-esteem, depression, behaviour and family functioning in sexually abused children. *Journal of Child Psychology and Psychiatry, 36*, 1077-1089.
- Sternberg, K., Lamb, M., Greenbaum, C., Cicchetti, D., Dawud, S., Cortes, R.M., Krispin, O., & Lorey, F. (1993). Effects of domestic violence on children's behavior problems and depression. *Developmental Psychology, 29*, 44-52.
- Tang, C.K. (1997). Psychological impact of wife abuse. (Experiences of Chinese women and their children). *Journal of Interpersonal Violence, 12*, 466-478.
- Timmons-Mitchell, J., Chandler-Holtz, D., & Semple, W.E. (1997). Post-traumatic stress disorder symptoms in child sexual abuse victims and their mothers. *Journal of Child Sexual Abuse, 6*, 1-14.
- Tong, L., Oates, K., & McDowell, M. (1987). Personality development following sexual abuse. *Child Abuse and Neglect, 11*, 371-383.
- Threlkeld, M.E., & Thyer, B.A. (1992). Sexual and physical abuse histories among child and adolescent psychiatric outpatients. *Journal of Traumatic Stress, 5*, 491-496.
- Trickett, P.K. (1993). Maladaptive development of school-aged, physically abused children: Relation-

- ships with the child-rearing context. *Journal of Family Psychology*, 71, 134–147.
- Walker, E., Downey, G., & Bergman, A. (1989). The effects of parental psychopathology and maltreatment on child behavior: A test of the diathesis-stress model. *Child Development*, 60, 15–24.
- Wallander, J.L., Varni, J.W., Babani, L., Banis, H.T., & Wilcox, K.T. (1988). Children with chronic physical disorders: Maternal reports of their psychological adjustment. *Journal of Pediatric Psychology*, 13, 197–212.
- Watts, W.D., & Ellis, A.M. (1993). Sexual abuse and drinking and drug use: Implications for prevention. *Journal of Drug Education*, 23, 183–200.
- Wells, R., McCann, J., Adams, J., Voris, J., & Dahl, B. (1997). A validation study of the structured interview of symptoms associated with sexual abuse (SASA) using three samples of sexually abused, allegedly abused, and nonabused boys. *Child Abuse and Neglect*, 21, 1159–1167.
- West, S., Sandler, I., Pillow, D., Baca, L., & Gerston, J. (1991). The use of structural equation modeling in generative research: Toward the design of a preventative intervention for bereaved children. *American Journal of Community Psychology*, 19, 459–480.
- White, S., Halpin, B.M., Strom, G.A., & Santilli, G. (1988). Behavioral comparisons of young sexually abused, neglected, and nonreferred children. *Journal of Clinical Child Psychology*, 17, 53–61.
- Williamson, J.M., Borduin, C.M., & Howe, B.A. (1991). The ecology of adolescent maltreatment: A multilevel examination of adolescent physical abuse, sexual abuse, and neglect. *Journal of Consulting and Clinical Psychology*, 59, 449–457.
- Wolfe, D., Sas, L., & Wekerle, C. (1994). Factors associated with the development of posttraumatic stress disorder among child victims of sexual abuse. *Child Abuse and Neglect*, 18, 37–50.
- Wolfe, V.V., Gentile, C., & Wolfe, D. (1989). The impact of sexual abuse on children: A PTSD formulation. *Behavior Therapy*, 20, 215–228.
- Young, R.E., Bergandi, T.A., & Titus, T.G. (1994). Comparison of the effects of sexual abuse on male and female latency-aged children. *Journal of Interpersonal Violence*, 9, 291–306.

Manuscript accepted 28 June 2002